



## Hawksbill Turtle (*Eretmochelys imbricata*)

Marine Turtle Specialist Group 2008 IUCN Red List status assessment.

**Assessors:** Jeanne A. Mortimer & Marydele Donnelly

The species account on the IUCN Red List web site documents the Rationale and supporting information for this Red List assessment (see below); that text is not repeated in this document.

### Location of supporting information text on the IUCN Red List:

<a href="#">Rationale</a> .....	see the <i>Justification</i> section in the species account on the IUCN Red List web site.
<a href="#">Range and Population</a> .....	see the <i>Geographic Range</i> section in the species account on the IUCN Red List web site.
<a href="#">Taxonomic Structure</a> .....	see the <i>Taxonomic Notes</i> section in the species account on the IUCN Red List web site.
<a href="#">Generation Length</a> .....	see the <i>Population</i> section in the species account on the IUCN Red List web site.
<a href="#">Nesting Population Size and Fecundity</a> .....	see the <i>Population</i> section in the species account on the IUCN Red List web site.
<a href="#">Habitats</a> .....	see the <i>Habitat and Ecology</i> section in the species account on the IUCN Red List web site.
<a href="#">Roles in the Ecosystem</a> .....	see the <i>Habitat and Ecology</i> section in the species account on the IUCN Red List web site.
<a href="#">Threats</a> .....	see the <i>Threats</i> section in the species account on the IUCN Red List web site.
<a href="#">Conservation Measures</a> .....	see the <i>Conservation Measures</i> section in the species account on the IUCN Red List web site.
<a href="#">Assessment Procedure</a> .....	see the <i>Justification</i> section in the species account on the IUCN Red List web site.
<a href="#">Population Trends and Conclusions</a> .....	see the <i>Population</i> section in the species account on the IUCN Red List web site.

Table of Contents	Page
<b>MTSG 2008 IUCN RED LIST STATUS ASSESSMENT</b>	
<a href="#"><u>Summary Information</u></a> .....	1
<a href="#"><u>Assessors</u></a> .....	1
<b><a href="#"><u>APPENDIX I: Tortoiseshell Trade Overview</u></a></b> .....	2
<a href="#"><u>History of the Trade</u></a> .....	2
<a href="#"><u>20<sup>th</sup> Century Trade</u></a> .....	2
<a href="#"><u>Hawksbills and CITES</u></a> .....	3
<a href="#"><u>The Japanese Tortoiseshell Trade</u></a> .....	4
<a href="#"><u>21st Century Global Trade</u></a> .....	5
<b><a href="#"><u>APPENDIX II: Regional Overviews</u></a></b> .....	7
<a href="#"><u>Indian Ocean</u></a> .....	7
<a href="#"><u>South Western Indian Ocean</u></a> .....	7
<a href="#"><u>North Western Indian Ocean</u></a> .....	8
<a href="#"><u>Central and Eastern Indian Ocean</u></a> .....	9
<a href="#"><u>Pacific Ocean</u></a> .....	10
<a href="#"><u>Western Pacific</u></a> .....	10
<a href="#"><u>Central Pacific</u></a> .....	13
<a href="#"><u>Eastern Pacific</u></a> .....	14
<a href="#"><u>Atlantic Ocean</u></a> .....	15
<a href="#"><u>Western Atlantic and Caribbean</u></a> .....	15
<a href="#"><u>Eastern Atlantic</u></a> .....	16
<b><a href="#"><u>LITERATURE CITED</u></a></b> .....	19
<b><a href="#"><u>ACKNOWLEDGEMENTS</u></a></b> .....	33

Table of Contents ( <i>continued</i> )	Page
Tables & Figures	
World	
<a href="#"><u>W-Figure 1</u></a> . World map with the geographic locations of the 25 Index Sites used for the 2006 MTSG Hawksbill Assessment.....	37
<a href="#"><u>W-Table 1</u></a> . Summary of estimated population change over 3 generations for 25 Index Sites based on Linear and Exponential extrapolation functions.....	38
Indian Ocean	
<a href="#"><u>IND-Table 1</u></a> . Indian Ocean localities of importance to <i>Eretmochelys imbricata</i> (n= 30), including 6 Index Sites for which quantitative data exist on past and present abundance.....	39
<a href="#"><u>IND-Table 2</u></a> . Quantitative evaluation of nesting activity and population trends in the Indian Ocean based on available <i>Past</i> and <i>Recent</i> estimates for <i>Eretmochelys imbricata</i> at 6 sites.....	41
<a href="#"><u>IND-Table 3</u></a> . Summary of estimated population change over 3 generations for the 6 Index Sites in the Indian Ocean.....	43
<a href="#"><u>IND-Table 4</u></a> . Changes in the size of Hawksbill populations in Seychelles during two decades (from the early 1980s to the early 2000s) at islands that have had different management regimes since the 1970s.....	47
<a href="#"><u>IND-Table 5</u></a> . Current population estimates and qualitative information about status and trends for reviewed Hawksbill populations in the Indian Ocean.....	48
Pacific Ocean	
<a href="#"><u>PAC-Table 1</u></a> . Pacific Ocean localities of importance to <i>Eretmochelys imbricata</i> (n= 19), including 7 Index Sites for which quantitative data exist on past and present abundance.....	61
<a href="#"><u>PAC-Table 2</u></a> . Quantitative evaluation of nesting activity and population trends in the Pacific Ocean based on available <i>Past</i> and <i>Recent</i> estimates for <i>Eretmochelys imbricata</i> at 7 sites.....	62
<a href="#"><u>PAC-Table 3</u></a> . Summary of estimated population change over 3 generations for the 7 Pacific Ocean Index Sites.....	65
<a href="#"><u>PAC-Table 4</u></a> . Detailed quantitative evaluation of nesting activity and population trends at 18 sites in Indonesia in the Western Pacific Ocean based on available <i>Past</i> and <i>Recent</i> estimates for <i>Eretmochelys imbricate</i> .....	69
<a href="#"><u>PAC-Table 5</u></a> . Current population estimates and qualitative information about status and trends for reviewed Hawksbill populations in the Pacific Ocean.....	73

Table of contents ( <i>continued</i> )	Page
Atlantic Ocean	
<a href="#"><u>ATL-Table 1</u></a> . Atlantic Ocean localities of importance to <i>Eretmochelys imbricata</i> (n= 34), including 12 Index Sites for which quantitative data exist on past and present abundance.....	82
<a href="#"><u>ATL-Table 2</u></a> . Quantitative evaluation of nesting activity and population trends in the Atlantic Ocean based on available <i>Past</i> and <i>Recent</i> estimates for <i>Eretmochelys imbricata</i> at 12 sites.....	84
<a href="#"><u>ATL-Table 3</u></a> . Summary of estimated population change over 3 generations for 6 Atlantic Ocean Index Sites for which there are historical data prior to the 1980s.....	89
<a href="#"><u>ATL-Table 4</u></a> . Summary of estimated population change over 3 generations for the Doce Leguas Cays of Cuba.....	93
<a href="#"><u>ATL-Table 5</u></a> . Estimated population change for 6 Atlantic Ocean Index Sites that have recorded increases in nesting populations since 1985, but for which no historical data exist prior to the 1980s.....	94
<a href="#"><u>ATL-Table 6</u></a> . Summary of estimated population change over 3 generations for 12 Atlantic Index Sites including: a) Caribbean sites with historic data prior to the 1980s; b) Caribbean sites lacking historic data prior to the 1980s; c) South Western & Eastern Atlantic sites; and d) Regional summaries of all index sites.....	96
<a href="#"><u>ATL-Table 7</u></a> . Current population estimates and qualitative information about status and trends for reviewed Hawksbill populations in the Atlantic Ocean.....	100
<a href="#"><u>ATL- Figure 1</u></a> . Recorded Hawksbill nesting in the Yucatan Peninsula 1977-2005.....	108

## Summary Information

**Class:** Reptilia; **Subclass:** Anapsida; **Order:** Testudines; **Family:** Cheloniidae; **Subfamily:** Chelonini

**Taxon name:** *Eretmochelys imbricata* (Linnaeus 1766)

**Common names:** Hawksbill turtle (English), tortue imbriquée (French), tortuga de carey (Spanish)

**Status:** *Critically Endangered* globally (CR A2bd)

**Distribution:** multiple genetic stocks occurring in tropical and subtropical waters around the world.

**Range:** Circumglobal, tropical to subtropical waters. Found in the waters of 108 countries, with nesting occurring in 70 countries.

**Habitats:** Adults nest on sandy beaches, primarily under vegetation. Post-hatchlings, small juveniles (<20-30 cm carapace length), and migrating animals are found in pelagic areas. Larger juveniles and adults forage in benthic habitats that include coral reefs and other hard bottom habitats, sea grass and algal beds, mangrove bays and creeks, and mud flats.

**Threats:** In the 19<sup>th</sup> and 20<sup>th</sup> centuries Hawksbill populations suffered dramatic declines in response to intense and prolonged exploitation of eggs and turtles for food and tortoiseshell. Today Hawksbills face not only purposeful exploitation, but also a suite of more insidious but equally destructive new threats. These include the loss of nesting and coral reef foraging habitat, incidental capture in fisheries, and marine and oil pollution.

## Assessors

Jeanne A. Mortimer<sup>1,2</sup>

<sup>1</sup> Department of Zoology, University of Florida, Gainesville, Florida 32611, USA

<sup>2</sup> Island Conservation Society, Victoria, Mahe, Seychelles. e-mail: [mortimer@ufl.edu](mailto:mortimer@ufl.edu)

Marydele Donnelly<sup>3</sup>

<sup>3</sup> Caribbean Conservation Corporation, Gainesville, Florida 32609, USA. e-mail: [marydele@cccturtle.org](mailto:marydele@cccturtle.org)

## **APPENDIX I: Tortoiseshell Trade Overview**

### **History of the Trade**

Tortoiseshell, the beautiful scutes of the carapace and plastron of the Hawksbill turtle, has been prized since ancient times. Surrounded by legend, tortoiseshell has been described as “one of the romantic articles of commerce, not only because of where it comes from, but because of the creatures from which it is obtained and the people engaged in the trade” (quoted in Parsons 1972). Jewelry and other tortoiseshell objects have been unearthed from pre-dynastic graves of the Nubian rulers of Egypt and excavated from the ruins of the Han Empire which ruled China in pre-Christian times. Over 2,000 years ago Julius Caesar considered the warehouses of Alexandria brimming with tortoiseshell to be the chief spoil of his triumph. By the early years of the 9<sup>th</sup> Century, caravans of Arab traders carried rhinoceros horn, ivory, and tortoiseshell throughout the Indian Ocean. For the next 1,000 years, the tortoiseshell trade flourished (Parsons 1972). Around 1700, during the Edo Period, the bekko (tortoiseshell) artisans of Japan established themselves at Nagasaki (Milliken and Tokunaga 1987).

The tortoiseshell trade has been closely linked to European discovery, conquest, and commerce around the world. The Portuguese, Dutch, French and English played major roles in the global trade; exploitation occurred throughout the world's tropical oceans and especially in the East Indies (i.e., modern day India, Indo China, Indonesia, Malaysia, and Philippines). The East Indies were a major source of the shell of antiquity, and these rich waters fittingly have been called the world's most productive seas for tortoiseshell (Parsons 1972). In the insular Pacific international trade did not develop until the mid 19<sup>th</sup> Century, but once established, it took a tremendous toll on the region's Hawksbills. For the next 150 years, tortoiseshell was a prized commodity in the Pacific, first with the sandal-wooders and then with the whalers (McKinnon 1975).

European Hawksbill fishing in the Caribbean began in the mid-17<sup>th</sup> Century and intensified throughout the 18<sup>th</sup> Century as demand increased (McClenachan *et al.* 2006). As they decimated local Hawksbill populations in one area after another, turtle fishermen moved from one site to the next. The plentiful Hawksbill resources of Central America were exploited for more than 100 years by traders, including Americans, who established the town of Bocas del Toro on the coast of Panama in 1826 (Parsons, 1972). Turtling was still a lucrative business in Cuba in 1885 when the village of Cocodrilos on the Isle of Pines was settled by turtle fishermen who emigrated from the Cayman Islands after its Hawksbills were gone (Carrillo *et al.* 1999). Over the next 100 years, many tens of thousands of Hawksbills were captured in the rich foraging grounds of the Cuban shelf.

### **20<sup>th</sup> Century Trade**

Tortoiseshell trade statistics are key to understanding the enormous and enduring effect that trade has had on Hawksbill populations around the world. In the early 20<sup>th</sup> Century, tortoiseshell was imported for luxury markets in Europe, the United States and Asia as the manufacture of combs and brushes, jewelry boxes, and tortoiseshell ornaments was “an established industry in almost every civilized country” (Seale 1917). Declines in Hawksbill populations were obvious in many areas by the first part of the century, as exemplified by expressions of “wanton destruction” in the Virgin Islands (Schmidt, 1916) and over exploitation in the Dutch East Indies (now Indonesia) (Dammerman 1929). Although existing records document an extensive trade in many countries, such as the 8,000 Hawksbills (8,000 kg) taken annually in the Philippines for the shell trade to Japan during World War I (Seale 1917) and 160,700 Hawksbills killed between 1918-1927 in the Dutch East Indies for export to Japan, Singapore and the Netherlands (Dammerman 1929), records for many other areas are incomplete.

During the 20<sup>th</sup> Century, Japan was the world's largest importer of tortoiseshell (Milliken and Tokunaga 1987, Groombridge and Luxmoore 1989). Although data are not available for imports in the first half of the century, Japanese statistics document the import of shell equivalent to more than 1.3 million large Hawksbills from around the world between 1950 and 1992, and more than 575,000 stuffed juveniles from Asia between 1970 and 1986 (Milliken and Tokunaga 1987, Groombridge and Luxmoore 1989).

Local trade in stuffed Hawksbills also flourished in the Indian Ocean, the Pacific and the Americas, especially in tourist areas. When Japanese, European, American and other Asian imports are considered along with the large quantities of tortoiseshell used locally in places like Sri Lanka and Madagascar, it is readily apparent that some millions of Hawksbills were killed for the tortoiseshell trade in the last 100 years.

## Hawksbills and CITES

In 1975, in recognition of its Endangered status, the Hawksbill was included on Appendices I (Atlantic population) and II (Pacific population) of CITES, the Convention on International Trade in Endangered Species of Wild Fauna and Flora, when the Convention came into force. By 1977 the entire species was moved to Appendix I to prohibit all international trade. Nevertheless, the global trade continued for a number of years, in large part driven by Japanese demand. At the end of 1992, Japanese imports ceased, but the industry continues to operate with stockpiled material.

- In the late 1970s more than 45 countries were involved in exporting and importing raw shell, with annual Japanese imports the equivalent of about 37,700 turtles (40,000 kg).
- Export and import levels remained exceptionally high until the mid-1980s as major trading nations slowly joined CITES. When they acceded to CITES in 1978, France and Italy took reservations (exceptions) to the Appendix I Hawksbill listing; these reservations were withdrawn in 1984 when they joined the EU.
- When Japan acceded to CITES in 1980, it also took a reservation on the Hawksbill and reduced its annual quota to the equivalent of 28,300 turtles (30,000 kg), based solely on the needs of its bekko industry.
- In 1985 CITES proposals by Indonesia and the Seychelles to place their Hawksbill populations on Appendix II to allow trade failed at the 5<sup>th</sup> CITES Conference of the Parties (COP 5). A similar Indonesian proposal at COP 6 in 1987 was withdrawn before the vote.
- A comprehensive report on the Japanese sea turtle trade by Milliken and Tokunaga in 1987 documented significant amounts of bekko trade with CITES countries. From 1980 to 1985, between 42% and 58% of all bekko imports originated in CITES countries, without proper export documents.
- In 1989 a detailed report commissioned by the CITES Secretariat found that Hawksbill populations were depleted or declining in 56 of 65 geopolitical units for which data were available and estimated that the annual global nesting population was a minimum of 15,000-25,000 Hawksbills. The authors concluded that the listing of the species on Appendix I was “unquestionably appropriate and must be maintained” (Groombridge and Luxmoore, 1989).
- On 1 April 1990, Japan reduced its annual bekko quota to the equivalent of 18,870 turtles (20,000 kg). In 1991, in an effort to avoid a U.S. embargo of its fish and fishery products, Japan agreed to further reduce its annual quota to the equivalent of 7,075 turtles (7,500 kg) by August 1991, to establish a zero quota on 1 January 1993, and to drop its Hawksbill reservation in July 1994. Japan also agreed to support the re-training of hundreds of bekko artisans. In the early 1990s, in response to the end of the Japanese trade, Cuba reduced its annual Hawksbill fishery quota from 5,000 turtles to 500.
- Since 1994, officials in Seychelles and Zanzibar have acquired tortoiseshell stocks from local artisans and subsequently burned them to demonstrate a commitment to ending the tortoiseshell trade (Khatib *et al.* 1996, Mortimer 1999). Cape Verde has shown similar commitment (Fretey *et al.* 2002).
- In 1997 and 2000, at CITES COP 10 and COP 11, Cuba proposed to sell its stockpiled tortoiseshell to Japan, and also proposed a continuation of the international trade in tortoiseshell taken from the 500 Hawksbills still captured each year. All these proposals failed.

- In response to regional disagreement generated by Cuban interest in moving Caribbean Hawksbills from Appendix I to II, the CITES Secretariat convened two regional Hawksbill dialogues in 2001 and 2002. The Dialogues encouraged regional cooperation by helping to establish Hawksbill priorities. As a result, resources for research, management and conservation have been generated.
- Although the tortoiseshell trade continues to threaten Hawksbills in numerous places, overall volume is substantially reduced. Thirty years after CITES came into force, the ban on international trade demonstrates its value over time in protecting Hawksbills. Above all, nesting increases in the Caribbean coincide with the enormous reduction in Hawksbill fishing in Cuban waters.
- In June 2007, Cuba informed CITES COP 14 that it would voluntarily institute a moratorium on its sea turtle fisheries in 2008. Although Cuba has a CITES Hawksbill reservation (exception) and reserves its right to dispose of its tortoiseshell stockpile, most nations are members of CITES and therefore cannot legally trade in tortoiseshell.

## The Japanese Tortoiseshell Trade

Twenty years ago, in their landmark report on Japan's sea turtle trade, Milliken and Tokunaga (1987) focused on providing estimates of the numbers of Hawksbills (and other species of sea turtles) represented by trade data so that the effect of Japanese exploitation around the world could be assessed. In particular, they cautioned that past exploitation is relevant to understanding and predicting current sea turtle population trends.

Estimates of the numbers of Hawksbills involved in the tortoiseshell trade are based on conversion factors calculated for each region by Milliken and Tokunaga (1987). Globally, the average Hawksbill produces 1.06 kg of tortoiseshell; but regionally, conversions are 0.74 kg in the Indian Ocean; 0.75 kg in Asia; 0.88 kg in Oceania; and 1.34 kg in the Caribbean. A combination of factors likely accounts for these differences, including regional variation in average adult size, as well as the relative proportion of adult and immature turtles represented in the trade. Some reports indicate that in the past the average turtle produced more shell than in recent decades. Adult turtles that survive long enough will continue to grow, so it follows that the average size of nesting animals tends to decline in an over-exploited population. In other cases, once nesting populations have been destroyed, hunters may shift their focus to foraging turtles which usually include immature animals. In the absence of specific historical information documenting the size classes of animals killed, the conversions we use in the present assessment are based on estimates provided by Milliken and Tokunaga. Based on the trade through 1992 (when legal Japanese imports ceased), the following information reveals the extent of the Japanese exploitation of global Hawksbill populations and the percent contribution of different regions to overall imports during 1950-1992.

- **Caribbean and Latin America (44.2%):** 29 countries provided the shell of 460,220 turtles (616,695 kg). Exports from Panama and Cuba were the equivalent of 152,070 and 106,948 turtles (203,774 kg and 170,047 kg, respectively), making them the most important sources of bekko in the world for Japan. Panama hosted the region's largest nesting Hawksbill assemblages until the latter part of the 20<sup>th</sup> Century. After 1961, Hawksbills in the Cuban trade were captured only at sea, but comprised adult and large immature animals.
- **Asia (20.8%):** 9 countries provided the shell of 387,020 turtles (290,265 kg). Exports from Indonesia were the equivalent of 155,654 turtles (116,741 kg), making it the most important source in the region and the third largest global supplier to Japan. Much of the shell exported from Singapore to Japan was probably of Indonesian origin (118,535 turtles, 88,901 kg). Asia was nearly the sole source of Japan's stuffed juvenile Hawksbill imports, as discussed below.
- **North America (15.1%):** the United States provided Japan with the shell of 199,490 turtles (211,463 kg) in two very large shipments, 142,241 kg in 1951 and 68,402 in 1954. The



countries of origin are unknown, but in all likelihood some quantity originated in U.S. Caribbean and Pacific territories.

- **Indian Ocean and East Africa (8.7%):** 15 countries provided the shell of 164,828 turtles (121,973 kg). Kenya and Tanzania, regional collection points, were the major exporters. Countries in the Northwestern Indian Ocean are notably absent from Japanese import statistics. As a non-CITES country, Maldives figured prominently in the trade after 1984 despite its national legislation protecting Hawksbills. Japanese imports therefore were in contravention of CITES Conf. Res. 4.25, which requires a nation with a reservation to treat an Appendix I species as Appendix II, with valid export documents from the country of origin.
- **Oceania (5.8%):** 6 countries provided the shell of 92,124 turtles (81,069 kg). A significant proportion of this trade is attributed to Australia until 1977 (29,109 turtles; 25,616 kg). Solomon Islands and Fiji were also important suppliers, especially in the final years of trading, with 40,982 and 14,490 turtles (36,064 and 12,751 kg, respectively). Fiji banned all tortoiseshell exports in January 1991 (Daly, 1991) but domestic tourist trade in Hawksbill curios and whole carapaces continues (PAC-Table 5).
- **Europe and West Africa (5.4%):** 10 countries provided the shell of 70,560 turtles (74,793 kg). The Netherlands was the largest exporter with the equivalent of 44,775 turtles (47,461 kg), but the source of this shell is unknown.
- In the 1970s, small lacquered Hawksbills became popular in Japan as symbols of long life. From 1970-1986 Japan imported 576,702 juvenile Hawksbills, mostly from Indonesia and Singapore but also from Taiwan, Province of China (32,075), the Ryukyus (13,438), Philippines (8698), Viet Nam (1195), Hong Kong (3549), and small quantities from a handful of other nations. Japan subsequently prohibited the trade, but continued to allow dealers to sell stocks acquired before July 1994. In December 1999, the dealers reported that they had a total of 135 stuffed sea turtles (TRAFFIC East Asia-Japan, 2000).
- Numerous irregularities in bekko imports occurred in the final years of Japan's trade under its CITES reservation. These included imports of shell from non-CITES countries that did not legally allow export of shell, as well as imports from countries known to have had too few turtles to supply the shell attributed to them. Based on these data, Japanese bekko imports from 11 of the 14 countries reported by the dealers in 1989 were illegal.
- The bekko stockpile in Japan includes raw shell and finished products. After Japan banned all imports in January 1993, annual Japanese domestic sales from stockpiled supplies remained high. Between July 1995 and July 1998 the stockpile was reduced from 188.4 to 102.73 tonnes (TRAFFIC East Asia-Japan, 2000). Information on subsequent annual sales and use is not available, but supplies would now be exhausted if utilization had continued at 28 tonnes a year after July 1998.
- Today, however, the bekko industry is intact, and Japanese consumer demand remains high. In January 2000, the valuable raw shell from abdominal plates ranged in price from JPY 30,000 per kg to JPY 150,000 per kg (US \$ 294-\$1470 at that time) (TRAFFIC East Asia-Japan, 2000).

## 21<sup>st</sup> Century Global Trade

Significant domestic trade in Hawksbill products is a major problem in many countries and, despite prohibitions on international trade and a reduction in its volume in the last decade, international and domestic trade remains an ongoing and pervasive threat in the Americas, Asia, and parts of Africa (Fleming 2001, Chacon 2002, TRAFFIC Southeast Asia 2004, van Dijk and Shepherd 2004, Brautigam and Eckert 2006, Reuter and Allan 2006).

- Some Japanese dealers have continued to import shell illegally as evidenced by numerous bekko shipments intercepted en route to or in Japan since the ban took effect (TRAFFIC East

Asia-Japan, 2000) and ongoing underground trade in southeast Asia to Japan and other destinations (van Dijk and Shepherd 2004, TRAFFIC Southeast Asia 2004).

- More than a decade after the Japanese prohibition on bekko imports took effect, van Dijk and Shepherd (2004) reported the interest of the Japan Bekko Association in acquiring Indonesia's remaining stockpiles of bekko.
- Although the volume of trade in Indonesia diminished significantly between 1991 and 2001, it is still substantial. The collection of tortoiseshell still occurs in numerous places, with most of the trade appearing to be disorganized and underground. Western Sumatra, Nias, and Papua are areas where significant exploitation and trade are known or suspected (van Dijk and Shepherd 2004).
- Those familiar with the trade warn that Indonesian stockpiles should be seized "as any indication of resumption of international trade of bekko could lead to requests from Indonesian traders to be allowed to sell their stockpiles" (van Dijk and Shepherd 2004).
- Surveys in Viet Nam in 2002 revealed an active international trade in tortoiseshell that had increased since 1999. Shell was purchased by tourists and foreigners buying in bulk for export to Hong Kong, Japan, South Korea, Taiwan (Province of China), Thailand, China and Asian communities in North America and Europe. Viet Nam subsequently instituted full protection for the Hawksbill (van Dijk and Shepherd 2004, TRAFFIC East Asia 2004).
- In recent reviews of the Lesser Antilles, Dominican Republic, Central America, Colombia and Venezuela, researchers provided evidence of extensive clandestine trade in sea turtles, including Hawksbills. Management and law enforcement are inadequate throughout the region (Brautigam and Eckert 2006, Reuter and Allan 2006).
- On 1 February 2007, the Kyodo News of Japan reported that Cuba would not seek to re-open the international tortoiseshell trade at the upcoming CITES meeting and noted Japanese disappointment given the long term support provided for the bekko industry. During 1991-2006, the Japanese government spent 735 million yen (US \$6M) for research on Hawksbill resources and 140 million yen (US \$1.1M) for projects to resume international trade, including trade with Cuba. The article also reported that the Ministry of Economy, Trade and Industry will support the bekko industry for another five years.

## **APPENDIX II: Regional Overviews**

### **Indian Ocean**

The coasts, islands and atolls of the Indian Ocean provide extensive nesting and foraging habitat for Hawksbills. This region has been an important center of the tortoiseshell trade since ancient times (Parsons 1972), and enormous quantities of shell were sold locally or exported to European and Asian markets during the 20<sup>th</sup> Century. Both historic and more recent exploitation have targeted vulnerable nesting Hawksbills and their eggs, with modern forms of transportation and navigational aids (i.e., GPS units) extending hunting forays to even the most remote areas. Juveniles were largely spared only until the second half of the 20<sup>th</sup> Century when the advent of the mask and snorkel, spear guns, and underwater torches facilitated the capture of turtles underwater (Mortimer 1984).

The full extent of population declines driven by the tortoiseshell trade may never be known, but, coupled with other major threats from direct exploitation (i.e., egg collection, hunting for meat, accidental capture in fisheries), destruction of nesting habitat (i.e., coastal development for tourism, human settlement, industry, etc.), and loss of marine foraging habitat due to pollution and other factors, most Hawksbill populations in the Indian Ocean are at risk. Exploitation for local consumption and tourist markets continues. Many populations are depleted, declining or remnants of larger assemblages. Extensive coral reef die-offs and subsequent reduction in foraging areas throughout the region have resulted from sea water warming events of 1998 and 2000 (Sheppard and Loughland 2002, Sheppard 2006); and these past events and the likelihood of more similar events in future are a cause for concern.

At many sites in the Indian Ocean region baseline surveys of Hawksbill populations were never conducted, but at certain other sites we have comparative data from surveys conducted both in recent years and 20-35 years ago. Another indicator of population trends is historic tortoiseshell trade statistics. Information derived from these sources indicate that within the last 135 years, (i.e., the time frame for this assessment in the Indian Ocean), nesting declines of significant proportions have occurred in Madagascar, Seychelles, Maldives, and probably India and Sri Lanka. The fate of smaller populations in places such as Egypt, Kenya, and Mozambique mirrors the demise of the larger aggregations. Madagascar's Hawksbills still may number about 1,000 females nesting annually, but this population is exploited and declining. The prognosis for the depleted nesting populations of Seychelles is good after 17 to 38 years of active protection on certain islands and 14 years of complete legal protection for turtles nationwide. Trends are unknown for two of the larger remaining Indian Ocean assemblages in Iran and Western Australia, but both populations face significant threats. Populations of hundreds of females nesting annually can still be found in the North Western Indian Ocean. IND-Table 1 provides an index of Hawksbill nesting sites in the Indian Ocean; qualitative and quantitative information is provided in IND Tables 2, 3, 4 and 5.

### **South Western Indian Ocean**

- Hawksbills have been hunted intensively for eggs, meat and shell in Mayotte, Mauritius, Kenya, Tanzania, and Mozambique; and at most of these sites nesting and foraging populations were still relatively abundant in the 1970s and 1980s (Hughes 1973, Groombridge and Luxmoore 1989, Frazier 1980). The exception is Mauritius, where the last known Hawksbill nesting attempt on the main island was in the mid-1970s (Mangar and Chapman 1996). In Mozambique coastal tribes have collected eggs intensely for decades (Groombridge and Luxmoore 1989). Today, although nesting female Hawksbills are rare in these countries; exploitation of eggs continues at most sites.
- Kenya and Tanzania were major suppliers of bekko to Japan in the second half of the 20<sup>th</sup> Century (IND Table 5). Zanzibar has served as a regional collection point for tortoiseshell since at least the 1890s (Frazier 1980). Coastal Bajunis in northern Kenya are among the world's

great turtle hunters (Frazier 1980). Foraging Hawksbills are still regularly encountered; but nesting animals have become rare.

- Formerly the site of one of the world's greatest concentrations of Hawksbills, Madagascar has a long history of tortoiseshell trade (Hughes 1973). Drastic declines in the early 20<sup>th</sup> Century are attributed to the killing of at least 1,600 adult turtles each year for more than 100 years. By the early 1970s, nearly 2,600 Hawksbills of all sizes were killed annually for the tourist trade (Hughes, 1973). Pressure on Hawksbills for meat, eggs, and shell remains intense (Ratsimbazafy 2004).
- Hawksbills were exploited in Seychelles ever since people first settled the islands in the late 17<sup>th</sup> Century, with trade intensifying in the 18<sup>th</sup> and 19<sup>th</sup> Centuries. By 1981, the long term trade in tortoiseshell had depleted the population (Mortimer 1984).
- In the 1970's, nesting Hawksbills in Seychelles received formal protection at several sites (i.e., Aride and Cousin islands, and Curieuse and the Ste. Anne Marine Parks). Informal protection was afforded Hawksbills at D'Arros/St. Joseph (Amirantes group) since the 1970s, and at the private islands of Bird and Cousine since 1992 (Mortimer 2004).
- In 1994 Seychelles Government enacted legislation protecting all species of sea turtles, and purchased and subsequently destroyed virtually all existing stocks of raw shell (Mortimer 1999). Today, the islands with the longest and most intense histories of protection boast increasing nesting populations; while sites where protection has been minimal or poaching ongoing, population declines continue (IND-Table 4). Hawksbill conservation in Seychelles is more advanced than in other parts of the Indian Ocean; nevertheless, coastal development threatens nesting habitat, and continued Government support is critical to the future of this globally significant population.

### ***North Western Indian Ocean***

The fact that Hawksbills in the north western Indian Ocean were only minimally involved in the Japanese tortoiseshell trade since at least 1950 probably explains the relative current abundance of Hawksbills in this region. Nevertheless, the species faces significant threats from entanglement in fishing gear, exploitation for meat and eggs, coastal development, and habitat degradation associated with oil production. Although traditions vary from country to country, in general where Muslims exploit sea turtles they are more likely to collect turtle eggs than to eat turtle meat; while foreign nationals working in the region consume both eggs and meat. Seismic exploration, pollution from spills and tanker washing, construction of port facilities, lights and vessel traffic have been recognized as major threats since the early 1980s (Ross and Barwani 1982, Frazier and Salas 1984, Miller 1989, Pilcher 1999). Oil pollution threatens nesting beaches and the region's extensive coral reef habitat. Corals have also been seriously impacted by warm water die-offs in 1998 and 2000.

- Hawksbill nesting sites in the Persian/Arabian Gulf include mainland and island beaches in Iran, eastern Saudi Arabia, the United Arab Emirates (UAE), Qatar, Kuwait and possibly Bahrain; but apparently not the estuarine coast of Iraq (Ross and Barwani 1982).
- Oman hosts significant Hawksbill nesting in the Gulf of Oman; while in the Gulf of Aden and the Arabian Sea, nesting occurs on the beaches of Oman, Yemen, Somalia, and Djibouti.
- In the Red Sea Hawksbills nest in Somalia, Sudan, Egypt, western Saudi Arabia, and Yemen.
- Vegetation loss and erosion caused by four-wheel drive vehicles threaten accessible nesting beaches in Oman, the UAE and probably most Arabian countries (Baldwin and Al-Kiyumi 1997).
- In the 1970s and early 1980s, Iran's nesting population was considered substantial, numbering perhaps 1,000 females (Kinunen and Walczak 1971, Ross and Barwani 1982); threats included egg predation by foxes and feral dogs and some incidental capture (Groombridge and Luxmoore 1989). In recent years, the population still appears to be large but faces very

significant threats of egg collection and predation, killing of nesting females, and incidental capture (Mobaraki 2003, 2004a, 2004b; Mobaraki and Elmi 2005) and high levels of egg utilization for medicinal purposes and animal feed (A. Mobaraki *in litt.* to CTURTLE 11 October 2004).

- Tortoiseshell commerce supported sea-faring and trade in the Red Sea for at least 2,000 years. By the second half of the 20<sup>th</sup> Century, populations were much reduced (Hirth and Abdel Latif 1980). Egypt's small nesting population is declining; the status of nesters in western Saudi Arabia is unknown.
- Decades of war and political strife have prevented surveys in Somalia, Eritrea, Sudan, and Yemen; so, the status of these populations is unknown.
- Somalis collect turtle eggs. Hawksbill populations were very reduced by the late 1960s although they were still hunted for tortoiseshell and eaten by coastal Bajunis in the south (Groombridge and Luxmoore 1989).
- In the late 1970s, hundreds of nesting Hawksbills were reported from Sudan's Suakin Archipelago; only meat was highly valued (Moore and Balzarotti 1977).
- In the mid-1990s, after the war with Ethiopia, Hawksbills were hunted in Eritrea for meat and eggs on a subsistence basis; small and medium-sized carapaces were sold in tourist shops (Hillman and Gebremariam 1996).
- In the 1960s and 1970s several hundred females a year nested annually in Yemen (Ross and Barwani 1982). At that time, eggs were collected, and some local hunting occurred (Groombridge and Luxmoore 1989).
- Hawksbills were killed and eggs were collected in Djibouti in the early 1980s; carapaces for sale in tourist shops may have been acquired locally or imported. The government prohibited egg collection and the killing of turtles in 1986 but did not regulate the sale of turtle products (Groombridge and Luxmoore 1989).
- Hawksbills are subject to heavy incidental capture in shark nets and shrimp trawlers in the Red Sea (Gebremariam *et al.* 1998).

### **Central and Eastern Indian Ocean**

Current trends for important Hawksbill nesting populations in Western Australia and the British Indian Ocean Territory (BIOT) are unknown. A relatively large but declining population is found in the Maldives. Historically large populations in Sri Lanka and India are much reduced.

- Trend data are not available for the very large Western Australian population which may number 2,000 or more females nesting each year. This assemblage is threatened by industrial development and habitat alteration. Most importantly, these turtles may be heavily exploited outside Australian waters (Limpus 2002).
- Trend data are not available for the smaller population of 300-700 nesting Hawksbills in the BIOT. Turtles have been protected there since the military took over the islands in the early 1970s, but many beaches are threatened by erosion (Mortimer and Day 1999).
- Today, less than 800 Hawksbills are estimated to nest annually on the thousands of islands and atolls of the Maldives although the population is believed to have once been very substantial (Groombridge 1982). Maldives has exported quantities of tortoiseshell to Sri Lanka since at least the 1920s, but the current nesting decline is the direct result of long term egg collection and over-exploitation since the religious ban on eating turtle meat was lifted in 1947 (Zahir and Hafiz 1997). Turtles, but not their eggs, were protected by a 10-year moratorium during 1995-2005 (Zahir and Hafiz 1997). In 2006, egg protection was initiated on a few islands.
- In the mid-19<sup>th</sup> Century Hawksbills were so abundant at India's Andaman and Nicobar islands that Malays visited Treis Island for six to eight months each year to collect shell (Kar and

Bhaskar 1982). But by the late 1990s only an estimated 250 Hawksbills were still nesting annually in the Andaman and Nicobar Islands (Andrews *et al.* 2006).

- Smaller numbers of Hawksbills also nest in India's Lakshadweep Islands where in the early 20<sup>th</sup> Century Hawksbills were "exceedingly numerous" (Laidlaw 1903, cited in Groombridge and Luxmoore 1989). They were still considered common in the late 1970s, but were hunted heavily for mainland tortoiseshell markets (Bhaskar 1978, Frazier 1980).
- India's shell exports in the second half of the 20<sup>th</sup> Century were the equivalent of 75,503 Hawksbills (55,872 kg). A small part of this shell (5,822 kg) was shipped to Japan. In 1977, shortly before India's accession to CITES, 50,050 kg were exported to Kuwait (Groombridge and Luxmoore 1989).
- Nesting females have nearly disappeared from Sri Lanka where abundant numbers of Hawksbills supported the tortoiseshell trade for centuries. By the 1920s Sri Lanka was importing more tortoiseshell than it exported, and by 1939 Hawksbill populations were greatly depleted (Deraniyagala 1939).
- Hawksbills were uncommon in Sri Lanka in the 1970s, with 50,000 fishermen dependent on turtle fishing (Salm 1981, cited in Groombridge and Luxmoore 1989). Eggs were avidly collected.
- Years of civil strife in Sri Lanka have prevented assessments in the north and east, but the prognosis for nesting Hawksbills is poor everywhere. In the 1980s and 1990s, hatcheries set up to generate tourist income operated for the benefit of tourists and mishandled hatchlings by holding them and scheduling daytime releases. As a result, few Hawksbills during that period can be expected to survive to maturity and return to breed. TCP, Sri Lanka's Turtle Conservation Project, has corrected this problem. In 2007, TCP protected about 20 Hawksbill nests, an increase over recent years (L. Ekanayake *in litt.* to J. Mortimer and M. Donnelly 22 Apr 2007 and 24 Apr 2007).
- In the late 1980s and more recently, small Hawksbills have been captured in the Gulf of Mannar between India and Sri Lanka (Groombridge and Luxmoore 1989, T. Kapurusinghe *in litt.* to J.A. Mortimer 2006); the origins of these turtles are unknown.
- Melaka, on the west coast of Peninsular Malaysia, hosts the second largest nesting population in Malaysia, an estimated 50-85 females nesting annually, which is significantly threatened by a history of intense egg exploitation, entanglement of adults in fishing gear, and massive coastal development (Mortimer *et al.* 1993).

In 2001, under the auspices of the Convention on Migratory Species (CMS), a Memorandum of Understanding on the Conservation and Management of Marine Turtles and Their Habitats in the Indian Ocean and South-East Asia (IOSEA MoU) was concluded. To date, 26 countries have signed the agreement, and numerous conservation activities are underway. The IOSEA Secretariat and Parties have established priorities, supported research, and convened annual meetings. The Year of the Sea Turtle was celebrated in the Indian Ocean in 2006.

## **Pacific Ocean**

### ***Western Pacific***

While Hawksbills face intense and varied threats in the western Pacific region, past and ongoing egg exploitation is currently the most pernicious problem for sea turtles there. Egg collection is not unique to this part of the world, but eating sea turtle eggs is deeply rooted in the cultures of Southeast Asia. Over decades, collection often has approached 100%, a situation exacerbated by relatively recent human settlement of previously uninhabited coastlines. As a result, during the 20<sup>th</sup> Century, many populations of Hawksbills and other sea turtles plummeted in Thailand, Indonesia, Malaysia, Myanmar, Philippines and Cambodia (Groombridge and Luxmoore 1989).

This decline was enhanced by both past and current exploitation of Hawksbills for the tortoiseshell trade, by continued take for meat, accidental capture in fisheries, and destruction of nesting habitat by unregulated coastal development. Their migratory nature makes sea turtles a resource shared by the various nations in this region. Many of these nations have burgeoning human populations which provide an incentive to intensively exploit marine resources. Because this region was historically one of the world's most famous Hawksbill breeding and foraging areas, its declining and depleted Hawksbill populations represent a significant global loss.

- On the east coast of Peninsular Malaysia, Hawksbills nest primarily in the states of Terengganu, Pahang, and Johor. Between 1956-1978 overall sea turtle nesting declined 43% in this region (Siow and Moll 1982); in the late 1970s several hundred Hawksbill nests were produced annually in Terengganu at Pulau Redang, Tanjung Galiga on the mainland, and Tioman island off the Pahang-Johor border. During 1987-1996, the number of nests in Terengganu ranged from 12-72 nests annually (Chan and Liew 1999), and declined to an average of 18 per year by 2002 (Liew 2002). Surveys conducted in 1990 estimated 100-200 egg clutches laid annually in Johor (Mortimer 1991b), and fewer than 100 in Pahang (Mortimer 1991a), with nesting levels at both sites reported by local inhabitants to be much lower than in previous years. Local informants attributed declines to over-exploitation of eggs, capture of turtles in commercial fishing gear (especially trawl nets), and destruction of nesting beach by coastal development (Mortimer 1991a, b). No recent data are available from Johor and Pahang.
- Green Turtle and Hawksbill eggs were collected intensely on the three turtle islands of western Sarawak in the South China Sea, with reports of organized egg collection dating from the early 19<sup>th</sup> Century. By 1936 few eggs were allowed to hatch. Egg production of all species declined by ~90% from the late 1920s to the mid 1980s (Groombridge and Luxmoore 1989). Additional eggs were imported from Indonesia to meet demand in Sarawak (Schulz, 1987). In the early 1980s, low level Hawksbill nesting was reported on the Sarawak mainland (de Silva 1982).
- Concern about capture for tortoiseshell in Sabah was expressed in 1927 (de Silva 1982); temporary bans on the hunting of Hawksbills were instituted in the 1920s and 1930s to control exploitation (Groombridge and Luxmoore 1989). Nevertheless, by the late 1960s the species was severely threatened (de Silva 1982).
- The Turtle Islands of Sabah in the Sulu Sea were privately held, and egg concessions were leased, until 1972. Hawksbill nests represent about 13.4% of the total egg production of the three Malaysian islands, with Pulau Gulisaan the most important site (Groombridge and Luxmoore 1989). In 1972 Malaysia established the Turtle Islands National Park, and egg production has been more or less stable since the late 1970s (PAC Table 2).
- When egg collection was banned in the Turtle Islands National Park, the demand for eggs was filled from the Philippine side of the Turtle Islands (five main islands). Tortoiseshell and stuffed specimens taken by armed fishermen or pirates in the Philippines or the boundary areas were smuggled into Malaysia (de Silva 1982). In 1976 a trans-border marine park was proposed to address turtle exploitation in the region.
- Only 90 years ago the outlying islands of the Philippines in the Sulu Archipelago were famous for their Hawksbill resources (Seale 1917). An intense 20<sup>th</sup> Century trade in eggs and Hawksbills of all sizes decimated populations, including the slaughter of nesting turtles by occupying Japanese forces in the early 1940s (de Celis 1982, de Silva 1982, Groombridge and Luxmoore 1989). Today Hawksbills nest in only low densities throughout the Philippines (Palma 1994, 1997).
- After World War II, the effects of heavy Hawksbill exploitation in the Philippines were evident. In the 1960s and 1970s exploitation was excessive; by the late 1970s Hawksbills were greatly diminished (Groombridge and Luxmoore 1989).
- From 1950 until 1985 substantial quantities of bekko were exported from the Philippines to Japan, Okinawa, and Taiwan (Province of China). In the 1970s smaller quantities of shell also

were exported to the French Pacific Islands, Italy, Singapore, and the USA (Groombridge and Luxmoore 1989).

- In 1996 Malaysia and the Philippines established the world's first trans-boundary marine park, the Turtle Island Heritage Protected Area, to protect the Philippine - Sabah Turtle Islands. Populations of Green and Hawksbill Turtles on the Filipino side of the Turtle Islands have declined 82% since the mid-1950s due to long-term exploitation (Palma 1997).
- With its 13,500 islands and abundant reefs, Indonesia historically has been one of the world's most important countries for Hawksbills and a center for the tortoiseshell trade (Parsons 1972).
- In 1929, Dammerman cautioned that customary egg collection, even more than Hawksbill fishing, endangered the species and recommended limits on hunting and egg gathering. Nearly 60 years later, Schulz (1987) expressed similar concerns.
- Ongoing declines have been noted in all areas of Indonesia surveyed in the last several decades as a result of both egg collection and the killing of larger animals. On many beaches, declines have been drastic in the last 20 years and the species is approaching extirpation in numerous areas (Schulz 1984, 1987; Meylan and Donnelly 1999; PAC Table 4).
- In eastern Indonesia 13 expeditions to all known or reported major Hawksbill and Green Turtle beaches between 1988 and 1995 revealed either low-level nesting or no nesting at all. In one area, fishermen reported that both Hawksbills and Green Turtles were more common in the waters off Pulau Ndana prior to 1970 when thousands could be seen during peak mating (Kitchener 1996).
- Japan was Indonesia's major trading partner for shell and stuffed juvenile Hawksbills in the second half of the 20<sup>th</sup> century. Between 1950 and 1986, bekko exports to Japan included the shell of 155,655 adult turtles and 428,859 stuffed juveniles (worked bekko) and in all likelihood an additional 59,215 large turtles and 88,539 stuffed specimens from Singapore (Milliken and Tokunaga 1987, Groombridge and Luxmoore 1989). Much of this trade occurred after Indonesia acceded to CITES in 1979. Indonesian authorities issued permits for bekko exports until 1985 (Milliken and Tokunaga 1987).
- Indonesia also exported large quantities of shell to other countries in the region, including Taiwan (Province of China), Singapore, Korea, and Hong Kong (Groombridge and Luxmoore 1989).
- Today Indonesia's tortoiseshell trade continues, with Yogyakarta (Java) a potential center of this commerce. In 2001 the level of trade was significantly lower than in 1991. The authors of a recent trade review concluded that the government did not appear to want to re-open international trade but may be interested in selling existing tortoiseshell stockpiles in Ujung Pandang and Sulawesi. The Japanese Bekko Association is reputed to be the potential buyer of this shell (van Dijk and Shepherd 2004).
- Surveys on continuing trade and stockpiles are needed in Ujung Pandang and in areas where significant exploitation and trade are known or suspected in Western Sumatra, Nias, and Papua (van Dijk and Shepherd 2004).
- Hawksbills from other countries are at risk as they migrate into Indonesian waters. Capture here has been identified as a major threat to NE Australian populations and possibly to Western Australian turtles (Limpus 2002).
- The very large rookeries in NE Australia are found in Torres Strait and Arnhem Land where Aborigines and Torres Strait Islanders eat turtle eggs, and no protection is provided. Declines have been underway in the nesting population at Milman Island, an index nesting beach for this aggregation, since work began there in 1990 (Limpus 2002). This population is subject to heavy exploitation during migration beyond Australian waters (Limpus 1997, 2004).

Capture and sale of Hawksbills at sea has been identified as a big problem in southeast Asia (van Dijk and Shepherd 2004), as demonstrated by Malaysia's seizure of two Chinese boats off the coast of Sabah in March 2007. A total of 300 Green and Hawksbill Turtles were on board, most of which were



dead (MSNBC News Services, 30 March 2007). On 2 May 2007 another Chinese boat was seized off Kalimantan Indonesia with 397 dead turtles on board, including 296 Hawksbills ranging from 20 to 90 cm carapace length, all preserved with formalin as stuffed specimens ([/www.turtle-foundation.org](http://www.turtle-foundation.org)).

Regional initiatives that benefit Hawksbill populations include the establishment of: a) the ASEAN Regional Conservation Program on marine turtle research and conservation in Southeast Asia (Brunei, Indonesia, Malaysia, Philippines, Singapore, Thailand, and Vietnam) in the 1990s; and b) the Memorandum of Understanding on the Conservation and Management of Marine Turtles and Their Habitats in the Indian Ocean and South-East Asia (IOSEA MoU) in 2001. The Southeast Asian Fisheries Development Center (SEAFDEC) initiated a research program for stock enhancement of sea turtles in 2004.

### **Central Pacific**

Sea turtles have been revered and used traditionally by the people of the Central Pacific for millennia. Important sources of protein, turtles figure prominently in religious ceremonies, art and legend of this region. Tortoiseshell was traded extensively and used for fish hooks, bride money, jewelry and other ornaments. Most likely, sea turtle populations in the Central Pacific were already heavily exploited prior to contact with European cultures. During the 1800s, the tortoiseshell trade developed in association with whalers and traders of sandalwood, beche de mer, pearl shell and salt pork (McKinnon 1975). These crews bought tortoiseshell wherever it was available, stopping at innumerable atolls and islands populated for trading. Fiji, Solomons, the Carolines, the Marianas and the Marshalls provided shell for this trade (Parsons 1972).

The history of the trade in the Solomon Islands demonstrates its far-reaching consequences on traditional Central Pacific cultures. In exchange for tortoiseshell, the islanders acquired iron and iron tools that simplified boat building and other work. This new efficiency allowed for more leisure time and the production of a fierce tomahawk which in turn fostered aggression and brutal raids for more tortoiseshell. According to McKinnon (1975), turtle hunting and head hunting were closely linked in the Solomons from the mid 1700s to early 1900s.

Since World War II, traditional taboos limiting use of sea turtles have broken down -- for example, the need to obtain permission to hunt turtles from hereditary rights holders. The flourishing souvenir trade in the region in the second half of the 20<sup>th</sup> Century further depleted populations. Technological advances including SCUBA gear have facilitated the capture of Hawksbills, and large outboard motors allowed access to once remote areas. Declines in Hawksbill populations at the three major island groups of the South Pacific -- Polynesia, Melanesia and Micronesia -- were well underway two or more decades ago as a result of intense egg collection and hunting (Pritchard 1982a, b; Balazs 1983; Johannes 1986; Groombridge and Luxmoore 1989; NMFS and USFWS 1998).

- The Solomon Islands and Fiji (Melanesia) have figured prominently in the global tortoiseshell trade for several hundred years.
- Hawksbill population declines in the Solomon Islands were reported at the turn of the 20<sup>th</sup> Century (McKinnon 1975) and again in the 1970s; despite increasing demand and higher prices, available supplies of tortoiseshell diminished (Pritchard 1982a). Nevertheless, the 200-300 females nesting annually in the Solomon Islands comprise the largest remaining Hawksbill population in Melanesia (NMFS and USFWS 1998). This population has been declining, however, and until at least the late-1990s most females were not surviving to nest for more than one season (90% of the nesting turtles being first time breeders) (Broderick 1998, Meylan and Donnelly 1999). Protection at the Arnavon Islands Community Marine Conservation Area (ACMCA) may enhance the prospects for survival (Broderick and Pita 2005).
- For decades Fiji islanders have ignored laws to protect Hawksbills, with perhaps as many as 2,000 turtles taken on feeding grounds each year through 1994 (Limpus 1997). Nest predation by feral mongooses is also problematic (Pritchard 1982a).
- Tortoiseshell products have been sold to tourists in Fiji for decades; raw shell exports to Japan began in the early 1960s and continued for nearly 30 years (Groombridge and Luxmoore 1989,

Japanese Trade Statistics); exports were banned in 1991 (Daly 1991). Significant tourist trade in tortoiseshell continues today (K. Mackay, pers. comm. to J. Mortimer & M. Donnelly, Feb 2007).

- A decade ago, an estimated several hundred Hawksbill females nested annually throughout the 2,200 islands of Micronesia, an area that extends from north of the Equator, east of the Philippines and southwest of Hawaii (NMFS and USFWS 1998). The 20 Hawksbills estimated to nest annually in Palau may be the largest single population in Micronesia; but, their eggs are poached, and increasing human disruption is a threat.
- Today a very small but increasing population of Hawksbills nests annually in the main Hawaiian Islands, with the east coast of the island of Hawaii an important area. Not all known nesting beaches are used every year. Although Hawksbills do not nest in the NW Hawaiian Islands today, historical records indicate that they may have nested there in the past (USFWS and NMFS 1998).

Regional initiatives include *The Year of the Sea Turtle* campaign, organized by the South Pacific Regional Environment Program in 1995, which promoted conservation and raised awareness of the plight of sea turtles in the region. In May 2004, a workshop on Southwest Pacific Hawksbills was the focus of one of two workshops which comprised the Second Western Pacific Sea Turtle Cooperative Research and Management Workshop (Kinan 2005). This initiative sponsored by the Western Pacific Regional Fishery Management Council is expected to support Hawksbill conservation in the region.

### ***Eastern Pacific***

The demise of Eastern Pacific Hawksbills was tied closely to the tortoiseshell trade. Coastal Indians in the Gulf of California and along the coast of Mexico traded tortoiseshell with the Spaniards during the colonial era (Del Barco 1980 cited in Clifton *et al.* 1982, Hardy 1929 cited in Clifton *et al.* 1982). Hawksbills were likely found in some numbers along the eastern coast of the Pacific several hundred years ago but have become very rare (Clifton *et al.* 1982, Seminoff *et al.* 2003b). A compilation of historical eyewitness accounts about the richness and abundance of marine life in the Gulf of California in the 16<sup>th</sup> to 19<sup>th</sup> Centuries (Sáenz-Arroyo *et al.* 2006) provides invaluable information about marine resources and their use by explorers, buccaneers, and local Indians. The 18<sup>th</sup> Century diaries of missionaries and others indicate that the Tres Marias Islands may have been an important Hawksbill breeding area; Hawksbills in the Gulf of California were exploited for commercial shell and jewelry industries on the mainland. At the end of the 18<sup>th</sup> Century, Hawksbills were reported to be the “commonest gift and meal of Indians from some missions” [Diario de los Expediciones a las Californias de José Longinos (1792) cited in Sáenz-Arroyo *et al.* 2006]. Reports of great turtle abundance in the region, presumably of all species, continued into the 19<sup>th</sup> Century.

- In the latter half of the 20<sup>th</sup> Century older fishermen reported that Hawksbills were abundant in the Gulf of California until the 1950s but were nearly eliminated by the lucrative tortoiseshell trade. At that time a small boat of fisherman could capture 5-7 Hawksbills in a few hours of night work along the east coast of Baja California. The shell was sold to a local prison where inmates were famous for crafting jewelry and other ornaments (Clifton *et al.* 1982).
- In recent years immature Hawksbills stranded or captured in neritic habitats at several sites within the Gulf of California and on the Pacific coast of Baja California are thought to originate from the population that nests in low levels on the mainland in the states of Jalisco and Nayarit and on the Tres Marias Islands (Seminoff *et al.* 2003b).
- Further south, Hawksbill nesting was reported twenty-five years ago, but not in significant numbers, in El Salvador, and along the Pacific coasts of Honduras, Nicaragua, Costa Rica, and Panama.
- Nesting beaches in El Salvador have been destroyed for tourist development, but the region’s greatest threats have been extensive egg collection and incidental capture in shrimp trawls (Cornelius 1982). Hawksbill eggs have been collected along with the hundreds of thousands of

turtle eggs traded annually in the region. In 2007 researchers reported 72 Hawksbills nested on three Salvadoran beaches between August and November (C. Hasbun pers. comm. to M. Donnelly, Feb 2008).

- In the late 1980s Hawksbills were present in Colombia, with the major nesting beaches located from Guapi south to the border with Ecuador. They were hunted for tortoiseshell and may have been captured accidentally in a major shrimp fishery (Groombridge and Luxmoore 1989). Hawksbills are still captured for meat and shell (INVEMAR 2002; C. Ceballos, in litt. 22 Aug 2007).
- In Ecuador Hawksbills nested in small numbers along much of the coast 25 years ago but most commonly between Manta and Cojimies; they foraged in the Galápagos Islands but did not nest there. Although there was little evidence of exploitation, tortoiseshell was sold in mainland tourist areas. Some Hawksbills were taken in shrimp trawls (Green and Ortiz-Crespo 1982).
- Based on the presence of immature Hawksbills in the Gulf of California, researchers concluded that the area should be a priority for regional recovery efforts (Seminoff *et al.* 2003b).

## Atlantic Ocean

### ***Western Atlantic and Caribbean***

Hawksbill populations of the wider Caribbean have been exploited for hundreds of years for European tortoiseshell markets and more recently for the Japanese market. Eggs were also exploited for human consumption. Historical accounts document the rich Hawksbill resources of the region, including the Yucatan (renowned as the best Hawksbill fishing in the Americas in the mid-1600s), the Doce Leguas Keys of Cuba, and the Caribbean coast of Central America (Parsons 1972). Historic records document European fishing of Hawksbills as early as the mid-17<sup>th</sup> Century (Craton and Saunders 1992 cited in McClenachan *et al.* 2006 (Web only PDF)), and indicate that exploitation intensified throughout the 18<sup>th</sup> Century with increasing demand (Williams 1969 cited in McClenachan *et al.* 2006).

For several hundred years turtle fishers followed a repeating pattern that entailed: intense exploitation, eventual decimation of local turtle stocks, and the need for them to move to a new good site to initiate exploitation anew. Turtling was still a lucrative business in the late 1800s when the town of Cocodrilos on Cuba's Isle of Pines was established in 1885 by turtle fishermen. At the turn of the 20<sup>th</sup> Century, a British call for thoughtful international sea turtle management in the Caribbean was not heeded (Schmidt 1916), and trade levels remained high. At about the same time the advice of a Dutch researcher who proposed protecting declining populations of Green and Hawksbill Turtles and their nests during the breeding season was similarly ignored (Boeke 1907 translated by Swinkels 2006). Instead, Caribbean Hawksbills were heavily exploited for much of the 20<sup>th</sup> Century. During this time, the region supplied enormous quantities of tortoiseshell to world markets.

- A recent review of historical records concluded that numerous major Hawksbill beaches existed throughout in the Caribbean just several hundred years ago. The authors cautioned that the loss of such an important animal as the Hawksbill in marine ecosystems cannot be ignored (McClenachan *et al.* 2006).
- The Doce Leguas Keys of Cuba were among the region's earliest known commercial Hawksbill fishing grounds (Parsons 1972). Historical records indicate thousands of nesting females were captured in Cuba annually during the 19<sup>th</sup> and 20<sup>th</sup> Centuries (ATL Table 7). From 1935-1994, 168,000 Hawksbills were taken on Cuban foraging grounds (Carrillo *et al.* 1999).
- The Cayman Islands, Jamaica, and the Florida Keys (USA) were once renowned for their Hawksbill fishing (Parsons 1972).
- The Wider Caribbean was a major source of bekko from 1950 through 1992, with shell exports to Japan from 27 countries being equivalent to 440,267 turtles (616,695 kg in total, with 170,047 kg from Cuba).

- Exports to Japan included large shipments from countries that support relatively few Hawksbills today, including Panama, Haiti, Nicaragua, and Jamaica; Haiti and the Cayman Islands served as regional collecting points (Milliken and Tokunaga 1987, Groombridge and Luxmoore 1989).
- In 1989 an in-depth global survey of Hawksbill populations undertaken for the CITES Secretariat concluded that the species was very reduced in the Western Atlantic and Caribbean (Groombridge and Luxmoore 1989).
- Small numbers of Hawksbills nest on many islands throughout the region but to date, little information has been collected systematically.
- Reviews conducted by Meylan (1999, 2001, 2002) estimated that fewer than 5,000 Hawksbill females nest annually in the region, with nesting populations depleted or declining in the majority of jurisdictions for which data are available. Despite recent increases in some areas, fewer than 5,000 females still nest in the Wider Caribbean today.
- In 2006, the WIDECAST network began compiling nesting data, recorded as numbers of crawls, for all species of Caribbean sea turtles. Some of these data are included in ATL-Table 7, where we estimated numbers of egg clutches by dividing by 1.8 (based on Mortimer & Bresson, 1999).
- Long-term data sets for protected areas in Puerto Rico, USVI, Jumby Bay Antigua, Barbados, and Cuba demonstrate nesting increases that coincide in time with the significant reduction of the Cuban fishery from 5,000 to 500 Hawksbills a year (this has spared ~55,000 large Caribbean Hawksbills since the early 1990s).
- Because numerous genetic haplotypes are shared by Caribbean Hawksbills (Bass *et al.*, 1996; Bowen *et al.* 2007), the exact contribution of individual countries to regional foraging populations in Cuban waters and elsewhere cannot be determined.
- Throughout the region, Hawksbill nesting and foraging habitat has been lost to beach development, sand mining, lights, and pollution.
- Hawksbills are captured accidentally in a variety of fisheries, including gill nets and pot fisheries.
- Consumptive utilization of turtles, including Hawksbills, is widespread and continues in the Lesser Antilles, Dominican Republic, Venezuela, Colombia, and Central America (Braütigam and Eckert, 2006).
- Long term data sets from Brazil demonstrate a significant nesting increase due to the protection efforts done by Projeto TAMAR since 1980 (Marcovaldi *et al.*, in press).

Regional initiatives to conserve sea turtles include two Western Atlantic Sea Turtle Symposia (WATS) held in 1984 and 1987. Research and conservation supported by governments and NGOs have generated excellent results. From 1995-2000 Hawksbills benefited from a five-year moratorium on sea turtle fishing in the eastern Caribbean. The SPAW Protocol of the Cartagena Convention and the Inter American Convention for the Protection and Conservation of Sea Turtles (IAC) help to conserve and protect Hawksbills. In 2006 the IAC adopted a resolution urging Parties to cooperate in supporting research and monitoring and addressing threats to Hawksbills from fisheries, illegal trade, and habitat destruction (IAC COP 3, 2006).

### ***Eastern Atlantic***

Sea turtles in the Eastern Atlantic have received little attention until recently. In the last several decades Hawksbills have been sighted or captured along the entire seaboard of the Eastern Atlantic from Western Sahara into the waters of Namibia. Nesting has been confirmed in some but not all of these countries (Brongersma 1982, Groombridge and Luxmoore 1989, Fretey *et al.* 2002). Today, Hawksbills are known to forage and nest in two areas, from Mauritania to west of the Ivory Coast, including Cape Verde, and in the Gulf of Guinea (Fretey *et al.* 2002).

Fewer than 100 Hawksbills now nest in all of West and Central Africa each year, with the best nesting on Bioko Island (Equatorial Guinea) and the islands of São Tomé and Príncipe. Subject to long-term exploitation, these populations are declining. Nesting is sporadic in other countries (Fretey 1998, Fretey *et al.* 2002), and historical accounts are limited, but Hawksbills may have nested in numerous places along the coast in the years before and after 1900 (Brongersma 1982, Groombridge and Luxmoore 1989). While Eastern Atlantic Hawksbill populations were depleted before baseline surveys were conducted, conservation activities currently underway for other species allow the collection of information on the region's remaining Hawksbills.

- Hawksbills have been observed in the waters of Western Sahara but not onshore, despite the availability of extensive nesting habitat.
- More than 20 years ago fishermen reported nesting along the southern half of Mauritania's remote coast to the border with Senegal (Groombridge and Luxmoore 1989). In Senegal about 10% of the turtles taken by fishermen are Hawksbills; the species may nest here as well. In Mauritania and Senegal drought increased the demand for meat and pressure on sea turtles (Groombridge and Luxmoore 1989).
- Sea turtle exploitation in Cape Verde dates to the mid 15<sup>th</sup> Century. In the late 1970s some hundreds of Hawksbills were taken annually for meat and shell, and eggs were collected daily (Groombridge and Luxmoore 1989). Prior to CITES prohibitions, tortoiseshell was shipped regularly to the Netherlands, and carapaces and shell were exported to Belgium. From 1996-1998 juvenile Hawksbills, but not nesting turtles, were sighted around five islands (López-Jurado *et al.* 2000).
- Cape Verde has been the only source of Japanese bekko in the region; exports from 1976-1983 were the equivalent of 432 turtles (458 kg).
- Baseline surveys in The Gambia found no evidence of Hawksbill nesting, but did record strandings of dead immature Hawksbills (Barnett *et al.* 2004).
- In Guinea Bissau, Hawksbill tracks identified in the Meio Islands in the early 1990s were later determined to be the tracks of another species (Barbosa *et al.* 1998). Four Hawksbill nests a year have been recorded on the island of Adonga in the Bijagos (Fretey *et al.* 2002).
- Hawksbills have been subject to subsistence take in Guinea where they may have nested in numbers in the late 1970s (Groombridge and Luxmoore 1989).
- Hawksbills were common in Sierra Leone 50 years ago, especially in the Turtle Islands off Sherbo Island, and they were reported to nest at Sussex and Bonthe (Groombridge and Luxmoore 1989). In 1991 no turtles or nests were sighted in a survey of Sherbo Island and the Turtle Islands (Fretey and Malaussena 1991).
- Nesting was reported in Liberia in the late 1800s (Brongersma 1982); in the early 1900s Hawksbills were eaten there (Groombridge and Luxmoore 1989).
- Nesting has not been confirmed in Ivory Coast. In the late 1960s Hawksbills probably were captured by the turtle fishery that took hundreds of animals annually (Groombridge and Luxmoore 1989).
- Old references document the presence of Hawksbills in Ghana, Benin, Cameroon, Gabon and Togo but do not identify them as nesting areas (Brongersma 1982).
- No information is available on Hawksbill nesting in Nigeria.
- During the 20<sup>th</sup> Century, Hawksbills nested in significant numbers in the Gulf of Guinea on 1) Bioko Island, Equatorial Guinea and 2) on São Tomé and Príncipe, but current nesting is much reduced. In 1999, 77 Hawksbill nests were reported from all of the Gulf of Guinea (Fretey *et al.* 2002).
- In the mid-1980s, during peak nesting, Hawksbills comprised a significant portion of the 50-100 turtles captured nightly in southern Bioko. At that time, intense capture for meat, shell and

eggs, including Russian exploitation in the 1970s, had reduced the population from the 1940s (Groombridge and Luxmoore 1989).

- In the early 1990s shell exploitation was identified as the cause of severely depleted Hawksbill populations in Bioko, but no estimates of nesting numbers were provided (Castroviejo *et al.* 1994). Two comprehensive surveys in southern Bioko in the late 1990s documented very little nesting (Tomás *et al.* 2000). No information is available for the mainland (Groombridge and Luxmoore 1989).
- Hawksbills nested frequently in São Tomé and Príncipe at the end of the 19<sup>th</sup> Century on the islands of São Tomé and Rolas where they were taken for meat, shell and eggs (Greef 1884 as cited in Groombridge and Luxmoore 1989). In the mid-1990s Hawksbills nested on most sandy beaches in the north, east, and south, but nesting estimates were not provided. Threats included sand-mining and egg collection (Graff 1996).
- Intense exploitation for the local tortoiseshell trade has been reported for decades in São Tomé (Brongersma 1982, Graff 1996) as well as export to Angola (Groombridge and Luxmoore 1989). A project to retrain tortoiseshell artisans and purchase their stocks of shell for destruction is underway (Fretey *et al.* 2002).
- Hawksbills have not been reported from Zaire or Angola, but they appear to forage as far south as Namibia (Groombridge and Luxmoore 1989).

Ongoing sea turtle conservation work in the Eastern Atlantic will yield new information about the region's Hawksbills, such as the use of reefs in Cameroon and São Tomé (Fretey *et al.* 2002). In the last 15 years the recapture of immature Hawksbills tagged in Brazil (Marcovaldi and Filippini 1991, Bellini *et al.* 2000, Grossman *et al.* 2007, A. Grossman. *in litt.* to J. Mortimer 3 Jul 2007) raises questions about the significance of trans-Atlantic crossings. Large coastal areas are relatively undeveloped and could support nesting, but unsustainable Hawksbill consumption is driven by severe poverty in the region. Legislation, and the will to enforce it, are needed in most areas. Habitat is threatened by expanding oil exploration and drilling in the Gulf of Guinea; garbage pollution is a growing threat (Formia *et al.* 2003).

Biologists and conservationists from many countries provide hope for the region's sea turtles (Formia *et al.* 2003). In 1999, the urgent need for regional cooperation culminated in A Memorandum of Understanding Concerning Conservation Measures for Marine Turtles of the Atlantic Coast of Africa, developed under the auspices of the Convention on Migratory Species (CMS). To date, 22 countries have signed this historic agreement.

## LITERATURE CITED

- Abreu-Grobois, F.A., Guzman, V., Cuevas, E. and Alba Gamio, M. (Compiladores). 2005. Memorias del Taller. Rumbo a la COP 3: Diagnóstico del estado de la carey (*Eretmochelys imbricata*) en la Península de Yucatán y determinación de acciones estratégicas. SEMARNAT, CONANP, IFAW, PRONATURA Península de Yucatán, WWF, Defenders of Wildlife. xiv + 75 pp.
- Akil, Y., Abudl, W., Sofyan, H., Jamaluddin, Tanaka, S. and Suganuma, H. 2004. Breeding status and recovery of resource of the endangered Hawksbill turtle in Western Java Sea and Natuna Sea, Indonesia. *Annual Report of Pro Natura Fund*. 13: 133-148. [in Japanese]
- Amarasooriya, D. 1996. Turtle hatcheries: Is it an additional disaster for the turtle fauna of Sri Lanka? Abstract. Paper presented at International Conference on the Biology and Conservation of the Amphibians and Reptiles of South Asia. August 1996. Kandy, Sri Lanka
- Andrews, H.V., Krishnan, S. and Biswas, P. 2006. Distribution and status of marine turtles in the Andaman and Nicobar Islands. In: K. Shanker & B. Choudhury (eds), *Marine Turtles of India*. pp. 33-57. Universities Press. Hydesabad, India. (contact: "Kartik Shanker" [kartikshanker@vsnl.net](mailto:kartikshanker@vsnl.net))
- Aronne, M. 2000a. Anidación semiartificial para la conservación de tortuga marina carey (*Eretmochelys imbricata*) en el Area Protegida de Cayos Cochinos, del 18 Junio al 30 Octubre 2000. Fundacion Hondureza Para Los Arrecifes Corales. 12 pp.
- Aronne, M. 2000b. Observaciones preliminares de la población anidadora de tortuga marina Carey *Eretmochelys imbricata* en la Reserva Biológica de Cayos Cochinos, 1999. PROARCAS. 16 pp.
- Baillie, J. and Groombridge, B. 1996. 1996 IUCN Red List of Threatened Animals. Gland, Switzerland: IUCN. 368 pp.
- Balazs, G.H. 1983. Sea turtles and their traditional use in Tokelau. Atoll Research Bulletin No. 279. Smithsonian Institution
- Baldwin, R.M. and Al-Kiyumi, A. 1997. Marine Turtles of the Sultanate of Oman. National Report for the IUCN MTSG Northern Indian Ocean Sea Turtle Workshop and Strategic Planning Session, 13-18 January, 1997, Bhubaneswar, India. 18 p.
- Barbosa, C., Broderick, A., and Catry, P. 1998. Marine Turtles in the Orango National Park (Bijagós Archipelago, Guinea-Bissau). *Marine Turtle Newsletter*. 81:6-7.
- Barnett, L.K., Emms, C., Jallow, A., Cham, A.M. and Mortimer, J.A. 2004. The distribution and conservation status of marine turtles in The Gambia, West Africa: a first assessment. *Oryx*. 38(2): 203-208.
- Bass, A.L., Good, D.A., Bjorndal, K.A., Richardson, J.I., Hillis, Z.M., Horrocks, J.A. and Bowen, B.W. 1996. Testing models of female reproductive migratory behavior and population structure in the Caribbean hawksbill turtle, *Eretmochelys imbricata*, with mtDNA sequences. *Molecular Ecology*. 5:321-328.
- Batibasaga, A. 2002. Sea turtle status and conservation initiatives in Fiji. In: I. Kinan (ed.), Proceedings of the Western Pacific Sea Turtle Cooperative Research and Management Workshop. February 5-8, 2002, Honolulu, Hawaii, USA. pp.115-118. Honolulu, HI: Western Pacific Regional Fishery Management Council.
- Beggs, J.A. 2006. Status of hawksbill sea turtle (*Eretmochelys imbricata*) nesting in Barbados, West Indies. M. Phil. thesis, University of the West Indies, Barbados.
- Beggs, J. A., Horrocks, J.A. and Krueger, B.H. 2007. Increase in hawksbill sea turtle *Eretmochelys imbricata* nesting in Barbados, West Indies. *Endangered Species Research*. 3:159-168.
- Bellini, C., Sanches, T.M. and Formia, A. 2000. Hawksbill turtle tagged in Brazil captured in Gabon, Africa. *Marine Turtle Newsletter*. 87:11-12.
- Ben Mohadji, F., Zarcach, H. and Mbindo, C. 1996. The status of sea turtle conservation in the Comoros. In: S.L. Humphrey & R.V. Salm (eds), *Status of Sea Turtle Conservation in the*

- Western Indian Ocean. pp. 125-132. Nairobi: IUCN/UNEP, UNEP Regional Seas Reports and Studies No. 165.
- Bennett, J.W. 1843 [1984 Facsimile Edition]. Ceylon and Its Capabilities. Kalapaluwawa, Rajagiriya, Sri Lanka: Trumpet Publishers Ltd., 427 pp.
- Bhaskar, S. 1978. Marine turtles in India's Lakshadweep islands. MTN 8:5.
- Bjorndal, K.A. 1997. Foraging ecology and nutrition of sea turtles. In: P.L. Lutz & J.A. Musick (eds). *The Biology of Sea Turtles*. pp: 199-232. CRC Press, Boca Raton, Florida.
- Bjorndal, K.A., Bolten, A.B. and Chaloupka, M.Y. 2000. Green turtle somatic growth model: evidence for density dependence. *Ecological Applications*. 10 (1): 269-282.
- Bjorndal, K.A., Bolten, A.B. and Chaloupka, M.Y. 2005. Evaluating trends in abundance of immature green turtles, *Chelonia mydas*, in the greater Caribbean. *Ecological Applications*. 15:304-314.
- Bjorndal, K.A., Bolten, A.B. and Lagueux, C.J. 1993. Decline of the nesting population of hawksbill turtles at Tortuguero, Costa Rica. *Conservation Biology* 7(4): 925-927.
- Bjorndal, K.A., and Jackson, J.B.C. 2003. Role of sea turtles in marine ecosystems--reconstructing the past. In: P.L. Lutz, J.A. Musick & J. Wyneken (eds), *Biology of Sea Turtles, Vol. II*. pp. 259-273. Boca Raton: CRC Press.
- Bouchard, S.S. and Bjorndal, K.A. 2000. Sea turtles as biological transporters of nutrients and energy from marine to terrestrial ecosystems. *Ecology*. 81: 2305-2313.
- Boulon, R., Jr. 1983. Some notes on the population biology of green *Chelonia mydas* and hawksbill *Eretmochelys imbricata* turtles in the northern U.S. Virgin Islands; 1981-1983. Rept. to NMFS, Grant No. NA82-GA-A-00044, 18 pp.
- Boulon, R., Jr. 1994. Growth rates of wild juvenile hawksbill turtles, *Eretmochelys imbricata*, in St. Thomas, U.S. Virgin Islands. *Copeia*. 1994(3):811-814.
- Bourjea, J., Ciccione, S. and Ratsimbazafy, R. In press. Marine turtles surveys in Nosy Iranja Kely, north-western Madagascar. Western Indian Ocean - *Journal of Marine Science*.
- Bowen, B.W., Bass, A.L., Garcia-Rodriguez, A., Diez, C.E., van Dam, R., Bolten, A., Bjorndal, K.A., Miyamoto, M.M. and Ferl, R.J. 1996. Origin of hawksbill turtles in a Caribbean feeding area as indicated by genetic markers. *Ecological Applications*. 6(2):566-572.
- Bowen, B.W., Grant, W.S., Hillis-Starr, Z., Shaver, D.J., Bjorndal, K.A., Bolten, A.B. and Bass, A.L. 2007. Mixed Stock analysis reveals the migrations of juvenile hawksbill turtles (*Eretmochelys imbricata*) in the Caribbean Sea. *Molecular Ecology*. 16: 49-60.
- Bowen, B.W. and Karl, S.A. 1997. Population genetics, phylogeography, and molecular evolution. In: P.L. Lutz & J.A. Musick (eds), *The Biology of Sea Turtles*. pp. 29-50. Boca Raton: CRC Press.
- Bräutigam, A.L. and Eckert, K.E. 2006. *Turning the Tide: Exploitation, Trade, and Management of Marine Turtles in the Lesser Antilles*. Central America, Colombia, and Venezuela. TRAFFIC International, Cambridge, UK, 547 pp.
- Broderick, D. 1998. Subsistence harvesting of marine turtles in the Solomon Islands. In: S.P. Epperly & J. Braun (compilers). Proceedings of the Seventeenth Annual Sea Turtle Symposium. U.S. Dep. Commer. NOAA Tech. Memo. NMFS-SEFSC-415. pp. 15-19.
- Broderick D., Moritz, C., Miller, J.D., Guinea, M., Prince, R.J. and Limpus, C.J. 1994. Genetic studies of the hawksbill turtle *Eretmochelys imbricata*: evidence for multiple stocks in Australian waters. *Pacific Conservation Biology* 1:123-131.
- Broderick, D. and Pita, J. 2005. Hawksbill Turtles in the Solomon Islands. In: I. Kinen (ed), Proceedings of the Second Western Pacific Sea Turtle Cooperative Research and Management Workshop. Volume 1: West Pacific Leatherback and Southwest Pacific Hawksbill Sea Turtles. Honolulu: Western Pacific Regional Fishery Management Council, pp. 101-102.
- Brongersma, L.D. 1982. Marine turtles of the Eastern Atlantic Ocean. In: K. Bjorndal (ed.), *The Biology and Conservation of Sea Turtles*. pp. 407-416. Washington, D.C.: Smithsonian Institution Press.



- Carr, A.F. 1956. *The Windward Road*. New York: Alfred Knopf, 258 pp.
- Carr, A.F., Carr, M. and Meylan, A.B. 1978. The ecology and migrations of sea turtles. 7. The west Caribbean green turtle colony. *Bulletin American Museum Natural History*. 162(1): 1-46.
- Carr, A.F. and Meylan, A.B. 1980. Extinction or rescue for the hawksbill? *Oryx*. 15(5): 449-450.
- Carr, A.F., Meylan, A.B., Mortimer, J.A., Bjorndal, K.A. and Carr, T. 1982. Preliminary survey of marine turtle populations and habitats in the western Atlantic. NOAA Tech. Memo. NMFS-SEFC-91, 91 pp.
- Carr, A.F. and Stanczyk, S. 1975. Observations on the ecology and survival outlook of the Hawksbill Turtle. *Biological Conservation*. 8: 161-172.
- Carrillo, E.C., Webb, G.J.W. and Manolis, S.C. 1999. Hawksbill turtles (*Eretmochelys imbricata*) in Cuba: an assessment of the historical harvest and its impacts. *Chelonian Conservation and Biology*. 3(2): 264-280.
- Castroviejo, J., Juste, J., Perez Del Val, J., Catelo, R. and Gil, R. 1994. Diversity and status of sea turtle species in the Gulf of Guinea islands. *Biodiversity and Conservation*. 3: 828-836.
- Ceballos-Fonseca, C. 2004. Distribucion de playas de anidacion y areas de alimentacion de tortugas marinas y sus amenazas en el Caribe Colombiano. *Bol. Invest. Mar. Cost.* 33: 77-99.
- Chacón, D. 2002. Diagnóstico sobre el comercio de las tortugas marinas y sus derivados en el istmo centroamericano. Red Regional para la Conservación de las Tortugas Marinas en Centroamérica (RCA). San José, Costa Rica. 247 pp.
- Chan, E-H. and Liew, H-C. 1999. Hawksbill turtles, *Eretmochelys imbricata*, nesting on Redang Island, Terengganu, Malaysia, from 1993 to 1997. *Chelonian Conservation and Biology*. 3(2): 326-329.
- Chaloupka, M.Y. and Limpus, C.J. 2001. Trends in the abundance of sea turtles resident in southern Great Barrier Reef waters. *Biological Conservation*. 102: 235-249.
- Chaloupka, M.Y. and Musick, J.A. 1997. Age, growth and population dynamics. In: P.L. Lutz & J.A. Musick (eds), *The Biology of Sea Turtles*. pp. 233-276. Boca Raton: CRC Press.
- Charuchinda, M. and Monanunsap, S. 1998. Monitoring survey on sea turtle nesting in the Inner Gulf of Thailand, 1994-1996. *Thai. Mar. Fish. Res. Bull.* 6: 17-25.
- Chevalier, J., Boitard, E., Bonbon, S., Boyer, J., Cuvillier, J.-M., Deproft, P., Dulorme, M., Giougou, F., Guyader, D., Lartiges, A., Leblond, G., Levesque, A., Lorvelec, O., Pavis-Bussi re, C., Rinaldi, C., Rinaldi, R., Roulet, M. and Thuaire, B. 2005. Update on the status of marine turtles in the Guadeloupean Archipelago (French West Indies). In: M.S. Coyne & R.D. Clark (compilers), *Proceedings of the Twenty-First Annual Symposium on Sea Turtle Biology and Conservation*. NOAA Tech. Memo. NMFS-SEFSC-528, pp. 135-136.
- Chevalier, J., Guyader, D., Boitard, E., Creantor, F., Delcroix, E., Deries, T., Deville, X., Guilloux, S., Nelson, L., Pavis, C., Roulet, M., Seman, J. and Thuaire, B. 2003. Discovery of an important hawksbill site in the Lesser Antilles: Trois Ilets Beach in Marie-Galante (Guadeloupe Archipelago - French West Indies). In: J.A. Seminoff (compiler), *Proceedings of the Twenty-Second Annual Symposium on Sea Turtle Biology and Conservation*. NOAA Tech. Memo. NMFS-SEFSC-503, p. 279.
- Cliffon, K., Cornejo, D.O. and Felger, R. S. 1982. Sea Turtles of the Pacific Coast of Mexico. In: Bjorndal, K. (ed.), *The Biology and Conservation of Sea Turtles*. Washington, D.C. Smithsonian Institution Press, pp. 199-209.
- Cornelius, S.E. 1982. Status of sea turtles along the Pacific Coast of Middle America. In: K. Bjorndal (ed.), *The Biology and Conservation of Sea Turtles*. pp. 211-219. Washington, D.C. Smithsonian Institution Press.
- Cordoba, J.A., Lopez, C.E. and Amorocho, D. 1998. Sea turtles in the Archipelago of San Andr s, Old Providence and Catleen-Caribbean, Columbia. In: S.P. Epperly & J. Braun (compilers). *Proceedings of the Seventeenth Annual Sea Turtle Symposium*. U.S. Dep. Commer. NOAA Tech. Memo. NMFS-SEFSC-415, p. 155.

- Cousin, J.-Y. 2001. Film: 1997-1998: Turtle's year in Mayotte. In: S. Ciccione, D. Roos & J.-Y. Le Gall (eds), *Proceedings of Connaissance et conservation des tortues marines du Sud-Ouest de l'océan Indien*. pp. 92-93. *Etudes et Colloques du CEDTM No 01*. March 2001.
- Cruz, G.A. and Espinal, M. 1987. National Report for Honduras. Western Atlantic Turtle Symposium II, Mayagüez, Puerto Rico, September 1987. Unpublished, 51 pp.
- Daly, T. 1991. Fiji bans export of turtle shell. *MTN* 52:1.
- Dammerman, K.W. 1929. Preservation of Wild Life and Nature Reserves in the Netherlands Indies. Fourth Pacific Science Congress. Java.
- Deraniyagala, P.E.P. 1939. The Tetrapod Reptiles of Ceylon, Volume 1. Testudines and Crocodilians. England: Dulau and Co., Ltd., 412 pp.
- de Celis, N.C. 1982. The status of research, exploitation and conservation of marine turtles in the Philippines. In: K. Bjorndal (ed.), *The Biology and Conservation of Sea Turtles*. Washington, D.C.: Smithsonian Institution Press, pp. 323-326.
- de Silva, G.S. 1982. The status of sea turtle populations in East Malaysia and the South China Sea. In: K. Bjorndal (ed.), *The Biology and Conservation of Sea Turtles*. pp. 327-337. Washington, D.C.: Smithsonian Institution Press.
- Diez, C.E. and van Dam, R.P. 2002. Habitat effect on hawksbill turtle growth rates on feeding grounds at Mona and Monito Islands, Puerto Rico. *Marine Ecology Progress Series*: 234: 301-309.
- Diez, C.E., van Dam, R.P. and Archibold, G. 2002. In-water survey of hawksbill turtles at Kuna Yala, Panamá. *Marine Turtle Newsletter* 96: 11-13.
- Dobbs, K.A., Miller, J.D., Limpus, C.J. and Landry, A.M., Jr. 1999. Hawksbill turtle, *Eretmochelys imbricata*, nesting at Milman Island, northern Great Barrier Reef, Australia. *Chelonian Conservation and Biology*. 3(2): 344-361.
- Domingue, J. and Mortimer, J.A. 2001. The impact of commercial fisheries on turtles in Seychelles. In: S. Ciccione, D. Roos & J.-Y. Le Gall (eds), *Proceedings of Connaissance et conservation des tortues marines du Sud-Ouest de l'océan Indien*. pp. 80-81. *Etudes et Colloques du CEDTM No 01*. March 2001. 135 pp.
- Dow, W.E. and Eckert, K.L. 2007. Sea Turtle Nesting Habitat – A Spatial Database for the Wider Caribbean Region. Wider Caribbean Sea Turtle Conservation Network (WIDECAST) and The Nature Conservancy. WIDECAST Technical Report No. 6. Beaufort, North Carolina.
- Eckert, K.L., Overing, J.A. and Lettsome, B.B. 1992. WIDECAST Sea Turtle Recovery Action Plan for the British Virgin Islands. Karen, K.L. (ed.). Kingston, Jamaica: UNEP Caribbean Environment Programme, CEP Tech. Rept. No. 15. 116 pp.
- Eckert, K.L. and Honebrink, T.D. 1992. WIDECAST Sea Turtle Recovery Action Plan for St. Kitts and Nevis. Eckert, K.L. (ed.). Kingston, Jamaica: UNEP Caribbean Environment Programme, CEP Tech. Rept. No. 17, 116 pp.
- Encalada, S.E., Lahanas, P.N., Bjorndal, K.A. Bolten, A.B. Miyamoto, M.M. and Bowen, B.W. 1996. Phylogeography and population structure of the Atlantic and Mediterranean green turtle *Chelonia mydas*: a mitochondrial DNA control region sequence assessment. *Molecular Ecology*. 5: 473-483.
- Fitzsimmons, N.N., Tucker, A.D. and Limpus, C.J. 1995. Long-term breeding histories of male green turtles and fidelity to a breeding ground. *Marine Turtle Newsletter* 68: 2-4.
- Fleming, E. 2001. *Swimming Against the Tide: recent surveys of exploitation, trade, and management of marine turtles in the Northern Caribbean*. TRAFFIC North America. 161 pp.
- Fletemeyer, J.R. 1984. National Report for Turks-Caicos. In: P. Bacon, F. Berry, K. Bjorndal, H. Hirth, L. Ogren & M. Weber (eds), *Proc. of the Western Atlantic Turtle Symposium*. Volume 3. pp. 409-422. Miami: RSMAS Printing.
- Formia, A., Tiwari, M., Fretey, J. and Billes, A. 2003. Sea Turtle Conservation Along the Atlantic Coast of Africa. *MTN* 100: 33-37.

- Frazier, J.G. 1980. Exploitation of marine turtles in the Indian Ocean. *Human Ecology* 8(4): 329-370.
- Frazier, J.G. 1982. Status of sea turtles in the Central Western Indian Ocean. In: K. Bjorndal (ed.), *The Biology and Conservation of Sea Turtles*. Washington, D.C. pp 385-389. Smithsonian Institution Press.
- Frazier, J.G. and Salas, S. 1984. The status of sea turtles in the Egyptian Red Sea. *Biological Conservation*. 30: 41-67.
- Frazier, J.G., Salas, S. and Didi, H. 2000. Marine Turtles in the Maldive Archipelago. *Maldives Marine Research Bulletin* 4. 80 pp.
- Fretey, J. 1998. Marine Turtles of the Atlantic Coast of Africa. UNEP/CMS Tech. Publ. No.1, 254 pp.
- Fretey, J. and Malaussena, J.-P. 1991. Sea turtle nesting in Sierra Leone, West Africa. *Marine Turtle Newsletter*. 54: 10-12.
- Fretey, J., Meylan, A. and Tiwari, M. 2002. The occurrence of the hawksbill turtle (*Eretmochelys imbricata*) in West Africa. In: A. Mosier, A. Foley & B. Brost (compilers). *Proceedings of the Twentieth Annual Symposium on Sea Turtle Biology and Conservation*. pp. 95-96. NOAA Tech. Memo. NMFS-SEFSC-477.
- Fuller, J.E., Eckert, K.L. and Richardson, J.I. 1992. WIDECASST Sea Turtle Recovery Action Plan for Antigua and Barbuda. CEP Tech. Rept. No. 16. Kingston, Jamaica: UNEP Caribbean Environment Programme, 88 pp.
- Garduño-Andrade, M., Guzmán, V., Miranda, E., Briseno-Duenas, R., and Abreu, A. 1999. Increases in hawksbill turtle (*Eretmochelys imbricata*) nestings in the Yucatán Peninsula, Mexico (1977-1996): data in support of successful conservation? *Chelonian Conservation and Biology* 3(2): 286-295.
- Gebremariam, T., Amer, A., Gebremariam, S. and Asfaw, M. 1998. Shrimp fishery in Eritrea: exploitation and legislation. In: G.M. Wamukoya & R.V. Salm (eds). *Report of the Western Indian Ocean Turtle Excluder Device (TED) Training Workshop*. Bandari College, Mombasa, Kenya. 27-31 Jan 1997. IUCN/EARO and IUCN/SSC MTSG, pp. 12-13.
- Graff, D. 1996. Sea turtle nesting and utilization survey in São Tomé. *MTN* 75:8-12.
- Grant, G.S., Craig, P. and Balazs, G.H. 1997. Notes on juvenile hawksbill and green turtles in the American Samoa. *Pacific Science* 51: 48-53.
- Gravier-Bonnet, N., Boulet, V., Bourjea, J., Ciccione, S. and Rolland, R. 2006. Mission Europa: 22 mai au 6 juin 2006. unpublished report. 19 pp.
- Grazette, S., Horrocks, J.A., Phillip, E. and Isaac, J. 2007. An assessment of the marine turtle fishery in Grenada, West Indies. *Oryx*. 41(3): 1-7.
- Green, D. and Ortiz-Crespo, F. 1982. Status of sea turtle populations in the Central Eastern Pacific. In: Bjorndal, K. (ed.), *The Biology and Conservation of Sea Turtles*. pp. 221-233. Washington, D.C.: Smithsonian Institution Press.
- Groombridge, B. 1982. The IUCN Amphibia-Reptilia Red Data Book, Part 1 Testudines, Crocodylia, *Rhynchocephalia*. IUCN, Gland, 426 pp.
- Groombridge, B. and Luxmoore, R. 1989. The green turtle and hawksbill (Reptilia: Cheloniidae): World status, exploitation, and trade. Lausanne, Switzerland: CITES Secretariat, 601 pp.
- Grossman, A., Bellini, C., Fallabrino, A., Formia, A., Mba Mba, J., Nzi Mba, J. and Obama, C. 2007. Second TAMAR-Tagged Hawksbill Recaptured in Corisco Bay, West Africa. *Marine Turtle Newsletter* 116: 26.
- Hamann, M., Chu The Cuong, Nguyen Duy Hong, Pham Thuoc, and Bui Thi Thuhien. 2006. Distribution and abundance of marine turtles in the Socialist Republic of Viet Nam. *Biodiversity and Conservation*. 25: 3703-3720.
- Hancock, J.M. 2007. Programa de conservación de tortugas marinas del caribe sur de Costa Rica Parque Nacional Cahuita. Temporada 2006. 65 pp.

- Harrison, E. Troëng, S. and Research Assistants. 2003. Report on the 2002 Green Turtle Program at Tortuguero, Costa Rica. Report to Caribbean Conservation Corporation and the Ministry of Environment and Energy of Costa Rica. 12 February 2003. 55 pages.  
<http://www.cccturtle.org/pdf/reports/TortugueroGreenTurtleReport2002>.
- Hewavisenthi, S. 1990. Exploitation of marine turtles in Sri Lanka: historic background and their present status. *Marine Turtle Newsletter*. 48: 14-19.
- Hill, M.S. 1998. Spongivory on Caribbean reefs releases corals from competition with sponges. *Oecologia*. 117: 143-150.
- Hillman, J.C. and Gebremariam, T. 1996. The status of sea turtle conservation in Eritrea. In: S.L. Humphrey & R.V. Salm (eds), *Status of Sea Turtle Conservation in the Western Indian Ocean*. pp. 41-56. Nairobi: IUCN/UNEP, UNEP Regional Seas Reports and Studies No. 165.
- Hirth, H. F. and Abdel Latif, E.M. 1980. A nesting colony of the hawksbill turtle *Eretmochelys imbricata* on Seil Ada Kebir Island, Suakin Archipelago, Sudan. *Biological Conservation*. 17: 125-130.
- Horrocks, J.A. 1992. WIDECAST Sea Turtle Recovery Action Plan for Barbados. Eckert, K.L. (ed.), CEP Tech. Rept. No. 12. Kingston, Jamaica: UNEP Caribbean Environment Programme, 61 pp.
- Howell, K. and Mbindo, C. 1996. The status of sea turtle conservation in Tanzania. In: S.L. Humphrey & R.V. Salm (eds.), *Status of Sea Turtle Conservation in the Western Indian Ocean*. pp. 73-80. Nairobi: IUCN/UNEP, UNEP Regional Seas Reports and Studies No. 165.
- Hughes, G.R. 1973. The survival situation of the hawksbill sea-turtle (*Eretmochelys imbricata*) in Madagascar. *Biological Conservation*. 5(2):114-118.
- Hunte, W. 1984. National Report for Barbados. In: P. Bacon, F. Berry, K. Bjorndal, H. Hirth, L. Ogren & M. Weber (eds.), *Proceedings of the Western Atlantic Turtle Symposium*. Volume 3. pp. 36-40. Miami: RSMAS Printing.
- INVEMAR. 2002. Determinación de la Distribución y del Estado Actual de Conservación de las Tortugas Marinas del Caribe Colombiano. Informe final Instituto de Investigaciones Marinas y Costeras (INVEMAR). Convenio SECAB-INVEMAR 052-029/01. 162 pp.
- IUCN. 2001a. *IUCN Red List Categories and Criteria: Version 3.1*. IUCN SSC. IUCN, Gland, Switzerland and Cambridge, UK. li + 30pp.
- IUCN. 2001b. IUCN Standards and Petitions Subcommittee. Red List Petitions Results. Species 36: 31-34.
- Johannes, R.E. 1986. A review of information on subsistence use of green and hawksbill sea turtles on islands under United States jurisdiction in the Western Pacific Ocean. NOAA-NMFS-SWFSC Admin. Rept. SWR-862, 41 pp.
- Joseph, D. 1984. National Report for Antigua and Barbuda. In: P. Bacon, F. Berry, K. Bjorndal, H. Hirth, L. Ogren & M. Weber (eds.), *Proceedings of the Western Atlantic Turtle Symposium*. Volume 3. Miami: RSMAS Printing.
- Kamel, S.J. and Mrosovsky, N. 2006. Deforestation: risk of sex ratio distortion in hawksbill sea turtles. *Ecological Applications*. 16(3): 923-931.
- Kapurusinghe, T. 2000. Community participation in turtle conservation in Sri Lanka. In: N. Pilcher & G. Ismail (eds.), *Sea Turtles of the Indo-Pacific: Research Management and Conservation*. pp. 35-44. Proceedings of the 2<sup>nd</sup> ASEAN Symposium and Workshop on Sea Turtle Biology and Conservation. ASEAN Academic Press Ltd.: London. 361 pp.
- Kar, C.S. and Bhaskar, S. 1982. Status of sea turtles in the Eastern Indian Ocean. In: K. Bjorndal (ed.), *The Biology and Conservation of Sea Turtles*. pp. 365-372. Washington, D.C.: Smithsonian Institution Press.
- Khatib, A.A., Khiari, S.K. and Mbindo, C. 1996. The status of sea turtle conservation in Zanzibar. In: S.L. Humphrey & R.V. Salm (eds.), *Status of Sea Turtle Conservation in the Western Indian Ocean*. pp. 81-88. Nairobi: IUCN/UNEP, UNEP Regional Seas Reports and Studies No. 165.

- Kiessling, I. 2003. Finding solutions: Derelict fishing gear and other marine debris in Northern Australia. A report for National Oceans Office and Department of the Environment and Heritage. 58 pp.  
<http://www.environment.gov.au/coasts/mbp/publications/north/pubs/marine-debris-report.pdf>
- Kinan, I. (ed.) 2005. Proceedings of the Second Western Pacific Sea Turtle Cooperative Research and Management Workshop. Volume 1: West Pacific Leatherback and Southwest Pacific Hawksbill Sea Turtles. 17-21 May, 2004. Honolulu, HI. Western Pacific Regional Fishery Management Council. Honolulu, HI USA.
- Kinunen, W. and Walczak, P.S. 1971. Persian Gulf sea turtle nesting surveys. Job Completion Rept., Division of Research Development, F-7-50, 16 pp.
- Kitchener, D.J. 1996. The status of green and hawksbill rookeries in Nusa Tenggara and Maluku Tenggara, Eastern Indonesia with observations on other marine turtles in the region. In: D.J. Kitchener & A. Suyanto (eds.), Proceedings of the 1st International Conference on Eastern Indonesian-Australian Vertebrate Fauna. Manado, Indonesia. November 22-26, 1994. pp. 97-109.
- La Gazette des Kares, 2006. Réseau tortues, Bulletin d'information mensuel. No. 2. Septembre 2006.
- Lagueux, C. 1998. Marine turtle fishery of Caribbean Nicaragua: human use patterns and harvest trends. Ph.D. Dissertation, University of Florida, Gainesville. 215 pp.
- Lagueux, C.J. and Campbell, C.L. 2005. Marine turtle nesting and conservation needs on the south-east coast of Nicaragua. *Oryx* 39(4): 398-405.
- Lagueux, C.J., Campbell, C.L. and McCoy, W.A. 2003. Nesting and conservation of the hawksbill turtle, *Eretmochelys imbricata*, in the Pearl Cays, Nicaragua. *Chelonian Conservation and Biology*. 4(3): 588-602.
- Lara-Ruiz, P., Lopez, P., Santos, G.G. Soares, F.R. 2006. Extensive hybridisation in hawksbill turtles (*Eretmochelys imbricata*) nesting in Brazil revealed by mtDNA analyses. *Conservation Genetics* 7(5): 773-891.
- Le Dien, D. and Broad, S. 1995. Exploitation of hawksbill turtles in Vietnam. *TRAFFIC Bulletin*. 15(2): 77-82.
- Lee, C. 2006. RM7mil home for turtles. New Straits Times. Local News. 9 Nov 2006.
- León, Y.M. and Bjorndal, K.A. 2002. Selective feeding in the hawksbill turtle, an important predator in coral reef ecosystems. *Mar Ecol-Prog Ser* 245: 249-258.
- Lescure, J. 1987. Tortues marines de l'Atlantique ouest. National Report for Martinique. Western Atlantic Turtle Symposium II, Mayagüez, Puerto Rico, September 1987. Unpublished, 27 pp.
- Liew, H-C. 2002. Status of marine turtle conservation research in Malaysia. In: I. Kinen (ed.), Proceedings of the Western Pacific Sea Turtle Cooperative Research and Management Workshop. pp. 51-56. Honolulu: Western Pacific Regional Fishery Management Council.
- Limpus, C.J. 1992. The hawksbill turtle, *Eretmochelys imbricata*, in Queensland: population structure within a southern Great Barrier Reef ground. *Wildlife Research*. 19: 489-506.
- Limpus, C.J. 1997. Marine turtle populations of Southeast Asia and the western Pacific Region: distribution and status. In: Y.R. Noor, I.R., Lubis, R., Ounsted, S., Troëng & A. Abdullah (eds.), Proceedings of the Workshop on Marine Turtle Research and Management in Indonesia. Bogor, Indonesia: Wetlands International, PHPA/Environment Australia, 197 pp.
- Limpus, C.J. 2002. Western Australia Marine Turtle Review. Unpublished report to Western Australian Department of Conservation and Land Management. 28 pp.
- Limpus, C.J. 2004. A Biological Review of Australian Marine Turtles. ii. Hawksbill Turtle, *Eretmochelys imbricata* (Linnaeus). Unpublished review produced by the Department of Environment and Heritage and Queensland Environmental Protection Agency. 58 pp.
- Limpus, C.J. and Miller, J.D. 2000. Final Report for Australian Hawksbill Turtle Population Dynamics Project. A Project Funded by the Japan Bekko Association to Queensland Parks and Wildlife

- Service. Prepared by Dr. Colin J. Limpus and Dr. Jeffrey D. Miller, Planning and Research Division, Queensland Parks and Wildlife Service, P.O. Box 155, Brisbane Albert Street, Qld 4002, Australia. 01 May 2000. Not for publication.
- Limpus, C.J., Miller, J.D., Guinea, M. and Whiting, S. 1997. Australian Hawksbill Turtle Population Dynamics Project. Paper presented at The International Workshop on the Management of Marine Turtles. 3-4 February 1997. Bali, Indonesia.
- Livingstone, S.R. 2006. Sea turtle ecology and conservation on the north coast of Trinidad. Chapter 5:83-113. PhD thesis. University of Glasgow, Scotland. Document not yet in the public domain. Contact: Suzanne R Livingstone SRLiving@odu.edu.
- López-Jurado, L.F., Cabrera, I., Cejudo, D., Evora, C. and Alfama, P. 2000. In: H.J. Kalb & T. Wibbels (compilers). Proceedings of the Nineteenth International Symposium on Sea Turtle Biology and Conservation. NOAA Tech. Memo. NMFS-SEFSC-443, pp. 245-247.
- Mangar, V. and Chapman, R. 1996. The status of sea turtle conservation in Mauritius. In: S.L. Humphrey & R.V. Salm (eds.), *Status of Sea Turtle Conservation in the Western Indian Ocean*. pp. 121-124. Nairobi: IUCN/UNEP, UNEP Regional Seas Reports and Studies No. 165.
- Marcovaldi, M.A. 2005. Brief overview of the hawksbill turtle in Brazil. In: I. Kinan (ed). Proceedings of the Second Western Pacific Sea Turtle Cooperative Research and Management Workshop. Volume 1: West Pacific Leatherback and Southwest Pacific Hawksbill Sea Turtles. Honolulu: Western Pacific Regional Fishery Management Council: pp. 91-95.
- Marcovaldi, M.A. and Filippini, A. 1991. Trans-Atlantic movement by a juvenile hawksbill turtle. *Marine Turtle Newsletter* 52:3.
- Marcovaldi, M.A., Lopez, G.G., Soares, L.S., Santos, A.J.B., Bellini, C. and Barata, P.C.R. in press. Fifteen years of hawksbill (*Eretmochelys imbricata*) sea turtle nesting in northern Brazil. *Chelonian Conservation and Biology*.
- Márquez M.R., Bravo G.R. and Castañeda A.P. 1987. Mexican Hawksbill Project, 1985-1986 Report. *Marine Turtle Newsletter* 41:3-5.
- McClenachan, L., Jackson, J. and Newman, M. 2006. Conservation implications of historic sea turtle nesting beach loss. *Frontiers in Ecology and the Environment* 4(6): 290-296.
- McGowan, A. Broderick, A.C., Frett, G., Gore, S., Hastings, M., Pickering, A., Wheatley, D., White, J., Witt, M.J. and Godley, B.J. MS in Review. Down but not out: Marine turtles of the British Virgin Islands.
- McIntosh, I., Goodman, K. and Parrish-Ballentine, A. 2003. Tagging and Nesting Research on Hawksbill Turtles (*Eretmochelys imbricata*) at Jumby Bay, Long Island, Antigua, West Indies. 2003 Annual Report. Prepared for the Jumby Bay Island Company, Ltd. WIDECAS. 21pp.
- McKeown, A. 1977. Marine turtles of the Solomon Islands. Honiara, Ministry of Natural Resources, Fisheries Division. 47 pp.
- McKinnon, J.M. 1975. Tomahawks, Turtles and Traders. A reconstruction in the circular causation of warfare in the New Georgian group. *Oceania*. 45(4): 290-307.
- Metcalf, J., K. Hampson, A. Andriamizava, R. Andrianirina, T. Cairnes, A. Gray, C. Ramiarisoa, and H. Sondotra. 2007. The importance of north-west Madagascar for marine turtle conservation. *Oryx* 41(2): 232-238.
- Meylan, A.B. 1982. Estimation of population size. In: K.A. Bjorndal (ed.), *Biology and Conservation of Sea Turtles*. pp. 135-138. Washington, D.C. Smithsonian Institution Press.
- Meylan, A.B. 1983. Marine Turtles of the Leeward Islands, Lesser Antilles. *Atoll Research Bulletin* 278: 1-43.
- Meylan, A. B. 1988. Spongivory in hawksbill turtles: a diet of glass. *Science*. 239: 393-395.
- Meylan, A.B. 1999. The status of the hawksbill turtle (*Eretmochelys imbricata*) in the Caribbean. Region. *Chelonian Conservation and Biology* 3(2): 177-184.



- Meylan, A. 2001. Global status review of the hawksbill turtle (*Eretmochelys imbricata*) with an emphasis on the wider Caribbean Sea. Paper presented at the First CITES Wider Caribbean Hawksbill Turtle Dialogue Meeting, 13-17 May 2001, Mexico City. 35 pp.
- Meylan, A.B. 2002. Global Status Review of the hawksbill turtle (*Eretmochelys imbricata*), with an emphasis on the wider Caribbean. An Addendum for the Second CITES wider Caribbean hawksbill turtle dialogue meeting, Cayman Islands (U.K.) 21-23 May 2002.
- Meylan, A., Castillo, I., Decastro Gonzalez, N., Ordoñez, C., Troëng, S., Ruiz, A. and Meylan, P. 2006. Bastimentos Island National Marine Park and Playa Chiriqui: Protected Areas Vital to the Recovery of the Hawksbill Turtle (*Eretmochelys imbricata*) in Caribbean Panama. Pp. 145-146. In: M. Frick, A. Panagopoulou, A.F. Rees & K. Williams (compilers). *Book of Abstracts. Twenty-Sixth Annual Symposium on Sea Turtle Biology and Conservation*. International Sea Turtle Society, Athens, Greece. 376 pp.
- Meylan, A.B. and Donnelly, M. 1999. Status justification for listing the hawksbill turtle (*Eretmochelys imbricata*) as Critically Endangered on the 1996 IUCN Red List of Threatened Animals. *Chelonian Conservation and Biology*. 3(2): 200-224.
- Meylan, A.B. and Redlow, A. 2006. *Eretmochelys imbricata* – Hawksbill Turtle. In: P.A. Meylan (ed.), *Biology and Conservation of Florida Turtles. Chelonian Research Monographs*. 3: 105-127.
- Miller, J. 1989. An assessment of the conservation status of marine turtles in the Kingdom of Saudi Arabia. Volume 1. Technical Report - Marine Turtles. Jeddah, Saudi Arabia: MEPA Report. No.9, 209 pp.
- Miller, J., Limpus, C.J. and Bell, I.P. 2000. Chapter 5. Nesting biology of *Eretmochelys imbricata* in the northern Great Barrier Reef. In: C.J. Limpus & J.D. Miller (compilers). Final Report for Australian Hawksbill Turtle Population Dynamics Project, funded by Japan Bekko Association, prepared for Queensland Parks and Wildlife Service. pp. 38-79.
- Milliken, T. and Tokunaga, H. 1987. The Japanese Sea Turtle Trade 1970-1986. A Special Report prepared by TRAFFIC (Japan). Washington DC: Center for Environmental Education, 171 pp.
- Mobaraki, A. 2003. Hawksbill nesting on Shidvar Island, Hormozgan Province, Islamic Republic of Iran. [http://www.ioseaturtles.org/archive\\_pom.php/year=2003](http://www.ioseaturtles.org/archive_pom.php/year=2003)
- Mobaraki, A. 2004a. Marine turtles in Iran: Results from 2002. *Marine Turtle Newsletter*. 104:13.
- Mobaraki, A. 2004b. Nesting of the Hawksbill Turtle at Shidvar Island, Hormozgan Province, Iran. *Marine Turtle Newsletter*. 103: 13.
- Mobaraki, A. and Elmi, A. M. 2005. Marine turtle tagging programme in the Islamic Republic of Iran. [http://www.ioseaturtles.org/archive\\_pom.php/year=2005](http://www.ioseaturtles.org/archive_pom.php/year=2005)
- Moncada, F. Carrillo, E., Saenz, A. and Nodarse, G. 1999. Reproduction and nesting of the hawksbill turtle (*Eretmochelys imbricata*) in the Cuban Archipelago. *Chelonian Conservation and Biology*. 3(2): 257-263.
- Monanunsap, S. 1997. Marine turtle research and management in Thailand. In: Y.R. Noor, I.R. Lubis, R. Ounsted, S. Troëng & A. Abdullah (eds). *Proceedings of the Workshop on Marine Turtle Research and Management in Indonesia*. Bogor, Indonesia: Wetlands International, PHPA/ Environment Australia, pp. 139-149.
- Moore, R.J. and Balzarotti, M.A. 1977. Report of 1976 Expedition to the Suakin Archipelago (Sudanese Red Sea). Results of marine turtle survey and notes on marine and bird life. Unpublished, 27 pp.
- Mortimer, J.A. 1984. Marine turtles in the Republic of the Seychelles: status and management. Report. on WWF project 1809, 1981-1984. Gland, Switzerland: IUCN/WWF, 80 pp.
- Mortimer, J.A. 1988. The pilot project to promote sea turtle conservation in southern Thailand with recommendations for a draft Marine Turtle Conservation Strategy for Thailand. Unpublished report to Wildlife Fund Thailand and the World Wildlife Fund US, 57 pp.

- Mortimer, J.A. 1991a. A Report on the Turtle Populations of the Islands of Pahang and Suggestions for their Management. Report submitted WWF/Malaysia, Jul 1991. 15 pp.
- Mortimer, J.A. 1991b. A Report on the Turtle Populations of the Islands off the East Coast of Johor and Suggestions for their Management. Report submitted to WWF/Malaysia, July 1991 (revision of Jan. 1990 report). 13 pp.
- Mortimer, J.A. 1991c. Recommendations for the Management of Turtle Islands Park, Sabah. Report submitted to WWF/Malaysia, Mar. 1991. 28 pp.
- Mortimer, J.A. 1995. Teaching critical concepts for the conservation of sea turtles. *Marine Turtle Newsletter*. 71: 1- 4.
- Mortimer, J. A. 1998. Turtle and Tortoise Conservation. Project J1, Environmental Management Plan of the Seychelles. Final report submitted to the Seychelles Ministry of Environment and the Global Environment Facility (GEF). Volume 1. 82 pp.
- Mortimer, J.A. 1999. World's first turtle shell stockpile to go up in flames as Miss World 1998 contestants look on. *Chelonian Conservation and Biology*. 3(2): 376-377.
- Mortimer, J.A. 2000. Conservation of hawksbill turtles (*Eretmochelys imbricata*) in the Republic of Seychelles. In: N. Pilcher & G. Ismail (eds), *Sea Turtles of the Indo-Pacific: Research Management and Conservation*. Proceedings of the 2<sup>nd</sup> ASEAN Symposium and Workshop on Sea Turtle Biology and Conservation. pp. 176-185. ASEAN Academic Press Ltd., London.
- Mortimer, J.A. 2002. Sea Turtle Conservation in the Arnavon Marine Conservation Area (AMCA): Observations and Management Recommendations made during a site visit in mid-2001. A report to The Nature Conservancy -- Asia Pacific Program. 68 pp.
- Mortimer, J.A. 2004. Seychelles Marine Ecosystem Management Project (SEYMEMP): Turtle Component. Final Report. Vol 1: Text, 243 pages. Vol 2: Appendix 1-11. 158 pp.
- Mortimer, J.A. 2006. Simple, yet Effective: Protection at the Nesting Beach. SWoT State of the World's Sea Turtles Report. 1: 8.
- Mortimer, J. and Bresson, R. 1999. Temporal distribution and periodicity in hawksbill turtles (*Eretmochelys imbricata*) nesting at Cousin Island, Republic of Seychelles, 1971-1997. *Chelonian Conservation and Biology*. 3(2): 318-325.
- Mortimer, J.A., Ahmad, Z. and Kaslan, S. 1993. The status of the hawksbill *Eretmochelys imbricata* and the green turtle *Chelonia mydas* of Melaka and Negeri Sembilan. *Malayan Nat. J.* 46: 243-253.
- Mortimer, J.A. and Day, M. 1999. Chapter 12: Sea turtle populations and habitats in the Chagos Archipelago. In: C.R.C. Sheppard & M.R.D. Seaward (eds). *Ecology of the Chagos Archipelago*. pp. 159-172. Linnean Society Occasional Publications 2.
- Mortimer, J.A., Day, M. and Broderick, D. 2002. Sea turtle populations of the Chagos Archipelago, British Indian Ocean Territory. In: A. Mosier, A. Foley & B. Brost (compilers), Proceedings of the Twentieth Annual Symposium on Sea Turtle Biology and Conservation. NOAA Tech. Memo. NMFS-SEFSC-477. pp. 47-49.
- Mortimer, J.A., Collie, J., Jupiter, T., Chapman, R., Liljevik, A. and Betsy, B. 2003. Growth rates of immature hawksbills (*Eretmochelys imbricata*) at Aldabra Atoll, Seychelles (Western Indian Ocean). P. In: J.A. Seminoff (compiler), Proceedings of the Twenty-Second Annual Symposium on Sea Turtle Biology and Conservation. NOAA Tech. Memo. NMFS-SEFSC-503. p. 247.
- Mortimer, J.A., Meylan, P.A. and Donnelly, M. 2007. Whose turtles are they, anyway? News and Views Perspective Article. *Molecular Ecology*. 16: 17-18.
- Musick, J.A. and Limpus, C.J. 1997. Habitat utilization and migration in juvenile sea turtles. In: P.L. Lutz & J.A. Musick (eds), *The Biology of Sea Turtles*. pp. 137-163. Boca Raton: CRC Press.
- Nava, M. and Uhr, A. 2007. 2006 Progress Report for Sea Turtle Conservation Bonaire. <http://www.bonairenature.com/turtles/PDF/STCB-Progress-Report-2006.pdf>.



- National Marine Fisheries Service and U.S. Fish and Wildlife Service. 1998. Recovery Plan for U.S. Pacific Populations of the Hawksbill Turtle (*Eretmochelys imbricata*). Silver Spring: National Marine Fisheries Service, 82 pp.
- Nietschmann, B. 1972. Hunting and fishing focus among Miskito Indians, eastern Nicaragua. *Human Ecology*. 1(1):41-67.
- Nietschmann, B. 1973. Between Land and Water: The Subsistence Ecology of the Miskito Indians. New York: Seminar Press, 279 pp.
- Nietschmann, B. 1981. Following the underwater trail of a vanishing species—the hawksbill turtle. *Nat. Geogr. Soc. Res. Rept.* 13:459-480.
- Okemwa, G.M., Simmons, N. and Mueni, E.M. 2004. The status and conservation of sea turtles in Kenya. *MTN* 105:1-6.
- Ottenwalder, J.A. 1981. Estudio Preliminar Sobre el Status, Distribucion y Biología Reproductiva de las Tortugas Marinas en la Republica Dominicana. Tesis para Optar por el Titulo de Licenciado en Biología. 113 pp.
- Ottenwalder, J.A. 1987. National Report for the Dominican Republic. Western Atlantic Turtle Symposium II, Mayagüez, Puerto Rico, September 1987 Unpublished, 52 pp.
- Palma, J.A.M. 1994. Marine turtle conservation in the Philippines. In: A. Nacu, R. Trono, J. Palma, D. Torres & F. Agas (eds). Proc. of the 1<sup>st</sup> ASEAN Symposium-Workshop on Marine Turtle Conservation, pp. 105-120. Manila, Philippines, 1993. Philippines: World Wildlife Fund.
- Palma, J.A.M. 1997. Marine turtle conservation in the Philippines and initiatives towards a regional management and conservation program. In: Y.R. Noor, I.R. Lubis, R. Ounsted, S. Troëng & A. Abdullah (eds). Proceedings of the Workshop on Marine Turtle Research and Management in Indonesia. Bogor, Indonesia: Wetlands International, PHPA/Environment Australia, pp. 121-138.
- Pandolfi, J.M., Bradbury, R.H., Sala, E., Hughes, T.P., Bjorndal, K.A., Cooke, R.G., McArdle, D., McClenachan, L., Newman, M.J.H., Paredes, G., Warner, R.R. and Jackson, J.B.C. 2003. Global trajectories of the long-term decline of coral reef ecosystems. *Science*. 301: 955-958.
- Parrish, A. and Goodman, K. 2006. Tagging and Nesting Research on Hawksbill Turtles (*Eretmochelys imbricata*) at Jumby Bay, Long Island, Antigua, West Indies. 2002 Annual Report. Prepared for the Jumby Bay Island Company, Ltd. WIDECAST. 22 p.
- Parsons, J.J. 1972. The hawksbill turtle and the tortoise shell trade. In: Études de géographie tropicale offertes a Pierre Gourou. Paris: Mouton, pp. 45-60.
- Pendoley, K.L. 2005. Sea turtles and the Environmental Management of Industrial Activities in North West Western Australia. Thesis presented for degree of Doctor of Philosophy, Murdoch University. 330 p. <http://wwwlib.murdoch.edu.au/adt/browse/view/adt-MU20060612.120104>
- Petro, G. 2002. Community empowerment: a case study: Wan Smolbag Turtle Conservation Program, Vanuatu. In: I. Kinan (ed.), Proceedings of the Western Pacific Sea Turtle Cooperative Research and Management Workshop. pp. 109-110. Western Pacific Regional Fisheries Management Council. Honolulu, HI.
- Pianka, E.R. 1974. Evolutionary Ecology. New York: Harper and Row, 356 pp.
- Pilcher, N.J. 2006. *Status of Sea Turtles in Qatar*. Supreme Council for the Environment and Natural Reserves, Doha, Qatar. 130 pp.
- Pilcher, N.J. 1999. The hawksbill turtle, *Eretmochelys imbricata*, in the Arabian Gulf. *Chelonian Conservation and Biology*. 3(2): 312-317.
- Pilcher, N.J. and Ali, L. 1999. Reproductive Biology of the hawksbill turtle, *Eretmochelys imbricata*, in Sabah, Malaysia. *Chelonian Conservation and Biology*. 3(2): 330-336.
- Polunin, N.V.C. and N.S. Nuijta. 1982. Sea turtle populations of Indonesia and Thailand. In: K.A. Bjorndal (ed.), *Biology and Conservation of Sea Turtles*. pp. 353-362. Washington, D.C.: Smithsonian Institution Press.

- Pritchard, P.C.H. 1982a. Marine turtles of the South Pacific. In: K. Bjorndal (ed.), *The Biology and Conservation of Sea Turtles*. pp. 253-262. Washington, DC: Smithsonian Institution Press.
- Pritchard, P.C.H. 1982b. Marine turtles of Micronesia. In: K. Bjorndal (ed.), *The Biology and Conservation of Sea Turtles*. pp. 263-274. Washington, DC: Smithsonian Institution Press.
- Quillard, M. 2001. Actions in Mayotte of the association Oulanga na Nyamba. In: S. Ciccione, D. Roos & J.-Y. Le Gall (eds), *Proceedings of Connaissance et conservation des tortues marines du Sud-Ouest de l'océan Indien*. pp. 94-95. Etudes et Colloques du CEDTM No 01. March 2001.
- Rader, H., Ela Mba, M.A., Morra, W. and Hearn, G. 2006. Marine turtles on the southern coast of Bioko Island (Gulf of Guinea, Africa), 2001-2005. *Marine Turtle Newsletter*. 111: 8-10.
- Rakotonirina, B. and Cooke, A. 1994. Sea turtles of Madagascar—their status, exploitation, and conservation. *Oryx*. 28 (1): 51-61.
- Ramohia, P. and Pita, J. 1996. Arnavon Islands Surveys. Unpublished Report to SPREP on the Regional Marine Turtle Conservation Programme (1995 Project) in the Solomon Islands. 20 p.
- Randriamiarana, H., Rakotonirina, B. and Maharavo, J. 1998. TED Experience in Madagascar. In: G.M. Wamukoya & R.V. Salm (eds), *Report of the Western Indian Ocean Turtle Excluder Device (TED) Training Workshop*. Bandari College, Mombasa, Kenya. 27-31 Jan 1997. pp. 16-17. IUCN/EARO and IUCN/SSC MTSG.
- Ratsimbazafy, R. 2004. Les tortues marines a Madagascar. Draft manuscript. 7 pp.
- Rees, A. F. and Papathanasopoulou, N. 2006. Masirah Island, Oman: Loggerhead turtles and much more under threat. [http://www.ioseaturtles.org/pom\\_detail.php?id=50](http://www.ioseaturtles.org/pom_detail.php?id=50)
- Reuter, A. and Allan, C. 2006. Tourists, Turtles and Trinkets: a look at the trade in marine turtle products in the Dominican Republic and Colombia. *TRAFFIC*. 12pp.
- Richardson, J.I., Bell, R. and Richardson, T.H. 1999. Population ecology and demographic implications drawn from an 11-year study of nesting hawksbill turtles, *Eretmochelys imbricata*, at Jumbay Bay, Long Island, Antigua, West Indies. *Chelonian Conservation and Biology*. 3(2): 244-250.
- Richardson, J.I., Hall, D.B., Mason, P.A., Andrews, K.M., Bjorkland, R., Cai, Y. and Bell, R. 2006. Eighteen years of saturation tagging data reveal a significant increase in nesting hawksbill sea turtles (*Eretmochelys imbricata*) on Long Island, Antigua. *Animal Conservation*. 9: 302-307.
- Ross, J.P. 1981. Hawksbill Turtle *Eretmochelys imbricata* in the Sultanate of Oman. *Biological Conservation*. 19: 99-106.
- Ross, J.P. and Barwani, M.A. 1982. Review of sea turtles in the Arabian area. In: K.A. Bjorndal (ed.), *The Biology and Conservation of Sea Turtles*. pp. 373-383. Washington, DC: Smithsonian Institution Press.
- Sáenz-Arroyo, A., Roberts, C. M., Torre, J., Cariño-Olvera, M., and Hawkins, J.P. 2006. The value of evidence about past abundance: marine fauna of the Gulf of California through the eyes of 16<sup>th</sup> to 19<sup>th</sup> century travelers. *FISH and FISHERIES* (7): 128-146.
- Salm, R.V. 1984. Sea Turtle Trade in Indonesia. IUCN/WWF Project 3108 Field Rept. No. 5, *Marine Conservation*. Bogor, Indonesia: IUCN/WWF, 50 pp.
- Salm, R.V., Jensen, R.A.C. and Papastavrou, V.A. 1993. *Marine Fauna of Oman: Cetaceans, Turtles, Seabirds and Shallow Water Corals*. Gland, Switzerland: IUCN, 66pp.
- Schulz, J.P. 1984. Turtle Conservation Strategy in Indonesia. IUCN/WWF Project No. 3108, Field Rept. No. 6. Bogor, Indonesia: IUCN/WWF.
- Schulz, J.P. 1987. Observations on sea turtles in Indonesia. Report to the IUCN Conservation Monitoring Center, 56 pp.
- Schmidt, J. 1916. Marking Experiments with Turtles in the Danish West Indies. Meddelelser Fra Kommissionen For Havundersogelser. Serie: Fiskeri. Bind V. Nr. 1. Kobenhavn. 26 pp.

- Seale, A. 1917. Sea products of Mindanao and Sulu, III. Sponges, tortoiseshell, corals and trepang. *Philippine Journal of Science*. 12: 191-211.
- Seminoff, J.A., Karl, S.A., Schwartz, T. and Resendiz, A. 2003a. Hybridization of the green turtle (*Chelonia mydas*) and hawksbill turtle (*Eretmochelys imbricata*) in the Pacific Ocean: indication of an absence of gender bias in the directionality of crosses. *Bulletin of Marine Science*. 73(3): 643-652.
- Seminoff, J.A., Nichols, W.J., Resendiz, A. and Brooks, L. 2003b. Occurrence of hawksbill turtles, *Eretmochelys imbricata* (*Reptilia Cheloniidae*) near the Baja California Peninsula, Mexico. *Pacific Science*. 57(1): 9-16.
- Sheppard, C.R.C. 2002. Island elevations, reef condition, and sea level rise in atolls of Chagos, British Indian Ocean Territory. *Cordio*. 3: 202-211.
- Sheppard, C. 2006. Longer-term impacts of climate change on coral reefs. In: I.M. Côté & J. Reynolds (eds), *Coral Reef Conservation*. pp. 264-290. Cambridge University Press.
- Sheppard, C. and Loughland, 2002. Coral mortality and recovery in response to increasing temperature in the southern Arabian Gulf. *Aquatic Ecosystem Health and Management*. 5(4): 395-402.
- Siow, K.T. and Moll, E. 1982. Status and conservation of estuarine and sea turtles in West Malaysian waters. In: K.A. Bjorndal (ed.), *The Biology and Conservation of Sea Turtles*. pp.339-347. Washington, DC: Smithsonian Institution Press.
- Smith, G.W., Eckert, K.L. and Gibson, J.P. 1992. WIDECAST Sea Turtle Recovery Action Plan for Belize. Eckert, K.L. (ed.). Kingston, Jamaica: UNEP Caribbean Environment Programme, CEP Tech. Rept. No. 18, 86 pp.
- Spring, S. 1982. Status of marine turtles in Papua New Guinea. In: K.A. Bjorndal (ed.), *The Biology and Conservation of Sea Turtles*. pp. 281- 289. Washington, DC: Smithsonian Institution Press.
- Stapleton, S. and Stapleton, C. 2004. Tagging and nesting research on hawksbill turtles (*Eretmochelys imbricata*) at Jumby Bay, Long Island, Antigua, West Indies: 2004 Annual Report. Prepared for the Jumby Bay Island Company, Ltd.. WIDECAST. 19 p.
- Stapleton, S. and Stapleton, C. 2006. Tagging and nesting research on hawksbill turtles (*Eretmochelys imbricata*) at Jumby Bay, Long Island, Antigua, West Indies: 2005 Annual Report. Prepared for the Jumby Bay Island Company, Ltd.. WIDECAST. 22 p.
- Suganuma, H. 2005. Hawksbill Sea Turtles in Indonesia. In: I. Kinan (ed.), *Proceedings of the Second Western Pacific Sea Turtle Cooperative Research and Management Workshop*. Volume 1: West Pacific Leatherback and Southwest Pacific Hawksbill Sea Turtles. 17-21 May 2004, Honolulu, HI. Western Pacific Regional Fishery Management Council: Honolulu, HI, USA, p. 103.
- Suganuma, H., Kamezaki, N. and Akil, Y. 1999. Current status of nesting populations of the hawksbill turtle (*Eretmochelys imbricata*) in the Java Sea, Indonesia. *Chelonian Conservation and Biology*. 3(2): 337-343.
- Swinkels, J. 2006. What we all should know...! Review on a 1904 Dutch fishery survey in the Caribbean Region. Unpublished MS. 6 pp.
- Tomás, J., Castroviejo, J. and Raga, J.A. 2000. Sea Turtles in the South of Bioko Island (Equatorial Guinea). In: H. Kalb & T. Wibbels (compilers). *Proceedings of the Nineteenth Annual Symposium on Sea Turtle Biology and Conservation*. pp. 247-250. U.S. Dept. Commerce. NOAA Tech. Memo. NMFS-SEFSC-443.
- TRAFFIC East Asia-Japan. 2000. Japan's Hawksbill Shell Trade Control System. A Briefing Prepared by TRAFFIC East Asia-Japan. 9 p.
- TRAFFIC Southeast Asia. 2004. The Trade in Marine Turtle Products in Viet Nam. Prepared for the Marine Turtle Conservation and Management Team by TRAFFIC Southeast Asia-Indochina. 48 pp.
- Troëng, S. 1996. Sea Turtle Conservation in Fiji. *Marine Turtle Newsletter* 73: 22-23.

- Troëng, S., Dutton, P.H. and Evans, D. 2005. Migration of hawksbill turtles *Eretmochelys imbricata* from Tortuguero, Costa Rica. *Ecography*. 28: 394-402.
- Tuato'o-Bartley, N., Morrell, T.E. and Craig, P. 1993. Status of sea turtles in American Samoa in 1991. *Pacific Science*. 47(3): 215-221.
- Valavi, H. 1999. Hawksbill nesting turtles in the Mound Islands. Unpublished report from Dept of Environment, Bushehr. 1 page abstract.
- van Dam, R.P. and Diez, C.E. 1997. Diving behavior of immature hawksbill turtles (*Eretmochelys imbricata*) in a Caribbean reef habitat. *Coral Reefs*. 16: 133-138.
- van Dijk, P.P. and Shepherd, C.R. 2004. Shelled out? A Snapshot of Bekko Trade in Selected Locations in Southeast Asia. TRAFFIC Southeast Asia, 29 pp.
- Vaughn, P.W. 1981. Marine turtles: a review of their status and management in the Solomon Islands. Unpublished report to WWF, 70 pp.
- Wamukoya, G.M. and Haller, R.D. 1995. The status and conservation of sea turtles in Kenya. In: J.A. Keinath, D.E. Barnard, J.A. Musick & B.A. Bell (eds), Proceedings of the Fifteenth Annual Workshop on Sea Turtle Biology and Conservation. pp. 336-339. NOAA Tech. Memo. NMFS-SEFSC-387.
- Wamukoya, G.M., Kaloki, F. and Mbindo, C. 1996. The status of sea turtle conservation in Kenya. In: S.L. Humphrey & R.V. Salm (eds), *Status of Sea Turtle Conservation in the Western Indian Ocean*. pp. 57-71. Nairobi: IUCN/UNEP, UNEP Regional Seas Reports and Studies No. 165.
- White, D. 2004. Marine Debris in Northern Territory waters 2003: WWF Report, WWF, Sydney. Online link: [http://wwf.org.au/publications/marine\\_debris\\_2003/](http://wwf.org.au/publications/marine_debris_2003/).
- Wetlands International, 1997. Noor, Y.R., Lubis, I.R., Ounsted, R. Troëng, S. and Abdullah, A. (eds). Proceedings of the Workshop on Marine Turtle Research and Management in Indonesia. Bogor, Indonesia: Wetlands International, PHPA/Environment Australia, 197 pp.
- Wilkinson, C.R. 2000. Status of Coral Reefs of the World: 2000. Global Coral Reef Monitoring Network. Australian Institute of Marine Science.
- Wilson, L., MacKay, K., Trevor, A. and Solomona, P. 2004. Melanesian Marine Turtles Conservation Forum. Workshop Report. 95 pp. <http://www.wwfpacific.org.fj>
- Witzell, W.N. 1983. Synopsis of biological data on the Hawksbill Turtle, *Eretmochelys imbricata* (Linnaeus, 1766). FAO Fisheries Synopsis No. 137, 78 p.
- Zahir, H. and Hafiz, A. 1997. Sea Turtles in the Maldives. National Report for the IUCN MTSG Northern Indian Ocean Sea Turtle Workshop and Strategic Planning Session, 13-18 January, 1997, Bhubaneswar, India. 16 pp.

## ACKNOWLEDGEMENTS

We thank the many people who assisted us during the preparation of this document. They made this in-depth assessment possible by compiling and sharing unpublished data, reviewing text, unearthing obscure documents, and providing excellent comments. They include:

Alberto Abreu Grobois	Karen Eckert	Loren McClenachan
Abdulrizak Orman Ali	Peter Eliazar	Anne Meylan
Monica Aureggi	Lalith Ekanayake	Jeffrey D. Miller
George Balazs	Angela Formia	Asghar Mobaraki
Rob Baldwin	Jack Frazier	Felix Moncado
Paul Basinthal	Jacques Fretey	Nick Pilcher
Ian Bell	Julie Garnier	John Pita
Claudio Bellini	David Godfrey	Earl Possardt
Rhema K. Bjorkland	Brendan Godley	Peter C.H. Pritchard
Karen Bjorndal	Hedelvy Guada	Ketut Putra
Alan Bolten	Vicente Guzman	Mireille Quillard
Rafe Boulon	Emma Harriso	Peter Ramohia
Jerome Bourjea	Carlos Hasbun	Bernhard Riegl
Damien Broderick	Zandy Hillis Starr	James I. Richardson
Cathi Campbell	Julia Horrocks	Rodney V. Salm
Claire Cayol	Brian Hutchinson	Jeff Seminoff
Milani Chaloupka	Thusan Kapurusinghe	Kartik Shanker
Eng Heng Chan	Rhema Kerr	Barbara Schroeder
Mickmin Charachinda	Barry Krueger	Isabel Marques da Silva
Stephane Ciccione	Cynthia Lagueux	Catherine Siota
Andrew Cooke	Min Min Lau	Luciano Soares
Alice Costa	Colin Limpus	Hiroyuki Suganuma
Eduardo Cuevas	Suzanne R. Livingstone	Jerome Swinkels
Gerry Davis	Sandy MacPherson	Manjula Tiwari
Carlos Diez	Kenneth T. MacKay	Sebastian Troëng
Kirstin Dobbs	Neca Marcovaldi	Robert van Dam
Jean-François Dontaine	Peri Mason	Rainer von Brandis
Wendy Dow	Rod Mast	Scott Whiting
	Andy McGowan	Hussein Zahir

We are especially grateful to the following persons and organizations for access to their unpublished information:

- Bahamas. for recent information and archival trade statistics:
  - Karen Bjorndal
  - Bernhard Riegl
  - Alan Bolten
  - Jeff Seminoff

- Barbados
  - Julia Horrocks
- Caribbean (general):
  - Loren McClenachan
  - Karen Eckert
  - The WIDECAST Network
- Cuba
  - Felix Moncado
- Indonesia:
  - Hiroyuki Suganuma - Everlasting Nature of Asia (ELNA). Special thanks for hours on the telephone providing volumes of information to JAM
  - Ketut Putra (Conservation International)
- Malaysia:
  - Paul Basinthal (Sabah Parks)
  - Min Min Lau (WWF Malaysia)
- Mexico, for unpublished long term trend data:
 

<ul style="list-style-type: none"> <li>◦ CONANP</li> <li>◦ Conanp-APFFLT</li> <li>◦ Desarrollo Ecologico Cd. del Carmen AC</li> <li>◦ Enlaces con tu Entorno AC</li> <li>◦ H. Ayuntamiento del Carmen</li> <li>◦ Marea Azul AC</li> <li>◦ Profepa</li> <li>◦ Pronatura Península de Yucatán</li> </ul>	<ul style="list-style-type: none"> <li>◦ Pronatura PPY</li> <li>◦ Quelonios AC</li> <li>◦ Secretaría de Ecología de Yucatán</li> <li>◦ Secretaria de Ecologia Gob. del Estado</li> <li>◦ SEMAR V Zona Naval</li> <li>◦ SEMARNAT</li> <li>◦ UNACAR</li> <li>◦ Universidad Autónoma de Campeche</li> </ul>
--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------
- Mozambique:
 

<ul style="list-style-type: none"> <li>◦ Alice Costa</li> <li>◦ Julie Garnier</li> </ul>	<ul style="list-style-type: none"> <li>◦ Isabel Marques da Silva</li> </ul>
------------------------------------------------------------------------------------------	-----------------------------------------------------------------------------
- Puerto Rico (Mona Is., Culebra Is, Caja de Muertos, & Humacao)
  - Carlos Diez (Chelonia, Inc.)
  - Robert van Dam (Chelonia, Inc.)

- Seychelles, for data used in the Mortimer 2004 report cited in this document:
  - Bird Island Lodge
  - Cousine Island Company
  - Denis Island
  - Fregate Island Private (FIP)
  - International Council for Bird Preservation (ICBP)
  - Island Conservation Society (ICS)
  - Marine Conservation Society Seychelles (MCSS)
  - Nature Protection Trust of Seychelles (NPTS)
  - Nature Seychelles
  - North Island Seychelles
  - Royal Society for Nature Conservation (RSNC)
  - Seychelles Centre for Marine Research and Technology - Marine Parks Authority (SCMRT-MPA)
  - Seychelles Islands Foundation (SIF)
  - Seychelles Ministry of Environment & Natural Resources (MENR)
- Solomon Islands: Arnavon Community Marine Conservation Area (ACMCA)
  - Peter Ramohia
  - John Pita
  - Catherine Siota
- Thailand: Gulf of Thailand & Andaman Sea
  - Monica Aureggi
  - Mickman Charuchinda
- U.S. Virgin Islands:
  - Zandy Hillis-Starr (National Park Service, Buck Island Reef NM)
  - Rafe Boulon
- Western & Central Pacific:
  - George Balazs (US NMFS)
  - Gerry Davis (US NMFS)
  - Barry Krueger (University of Barbados)
  - Kenneth.T. MacKay (University of the South Pacific)

We thank the MTSG Assessment Steering Committee (ASC) for its careful one-month-long review of the draft Assessment prior to its posting on the MTSG website for general review. The members of the MTSG-ASC include the following:

Milani Chaloupka (Chairman, MTSG-ASC)

Alan Bolten

Annette C. Broderick

Kirstin Dobbs

Peter Dutton

Colin Limpus

Jeffrey D. Miller

Jack Musick

Peter C. H. Pritchard

Kartik Shanker


Peter Paul van Dijk

After comments of the MTSG-ASC were incorporated, the draft document was then posted on the MTSG website for a period of two months to enable general comment and review by the wider MTSG membership and other interested parties. We are grateful to the many people who took the time to submit their comments about the draft document in particular and the IUCN Red Listing process in general. Some people engaged in the online discussion that ensued on the MTSG Listserv; some sent their comments directly to the assessors (JAM & MD) and MTSG-ASC Chair (MC); and some did both. All MTSG postings and comments received have been compiled for the record. Those who participated in this phase of the process include:

Alberto Abreu Grobois	Cathi Campbell	Marydele Donnelly
Diego Amorocho	Claudia Ceballos	Carlos Drews
George Balazs	Milani Chaloupka	Karen Eckert
Ana Barragan	B.C. Choudbury	Sheryan Epperly
Karen Bjorndal	Eduardo Cuervas	Angela Formia
Alan Bolten	Charlotte de Fontaubert	David Godfrey
Brian Bowen	Carlos Diez	Matthew Godfrey
Joaquin Buitrago	Kirstin Dobbs	Brendan Godley
Alice Grossman	Peter Meylan	Jeffrey Seminoff
Hedelvy Guada	Felix Moncado	Kartik Shanker
Vicente Guzman	Jeanne A. Mortimer	Jim Spotilla
Mark Hamann	Nicholas Mrosovsky	Todd Steiner
Emma Harrison	Wallace J. Nichols	Hiroyuki Suganuma
Selina Heppell	Ronald Orenstein	Jesus Tomas
Paul Hoetjes	Frank Paladino	Sebastian Troëng
Julia Horrocks	Kellie Pendoley	Robert van Dam
Brian Hutchinson	Nicholas Pilcher	Richard van der Wal
Barry Krueger	Pamela Plotkin	Edith van der Wal
Cynthia Lagueux	Earl Possardt	Marc Ward
Colin Limpus	Bob Prince	Grahame Webb
Sandy MacPherson	Anders Rhodin	Scott Whiting
Charlie Manolis	Marc Rice	Blair Witherington
Neca Marcovaldi	Laura Sarti	Zahirul Islam
Anne Meylan	Barbara Schroeder	

We are grateful for financial support provided to JAM by the IUCN Marine Turtle Specialist Group and the US Fish & Wildlife Service, and to MD by Caribbean Conservation Corporation. Special thanks for advice provided by Milani Chaloupka (Chair of the MTSG Assessment Steering Committee), Alberto Abreu Grobois (former Chair of the MTSG), and Jeffrey Seminoff (former MTSG Red List Focal Point); and for assistance from Rod Mast (Co-Chair MTSG), Brian Hutchinson, and Nick Pilcher (Co-Chair MTSG) during preparation of the document.





**W-Figure 1.** World map with the geographic locations of the 25 Index Sites used for the 2006 MTSG Hawksbill Assessment. For the rationale for inclusion of each location as an Index Site see: IND-Table 1, PAC-Table 1, and ATL-Table 1.

**W-Table 1.** Summary of estimated population change over 3 generations for 25 Index Sites based on Linear and Exponential extrapolation functions (IUCN, 2001a). The derivations of these figures are detailed in the following tables: for the Indian Ocean in IND-Table 3; for the Pacific Ocean in PAC-Table 3; and Atlantic Ocean in ATL-Table 6.

	Number of Index Sites	Raw Data & <u>Linear</u> Functions			Raw Data & <u>Exponential</u> Functions		
		3 Generations Back: in Indo-Pacific 1870); in Atlantic (1901)	2005	% Change over 3 Generations	3 Generations Back: in Indo-Pacific 1870); in Atlantic (1901)	2005	% Change over 3 Generations
Indian Ocean	6	30,430	1,893	-93.8 %	39,517	2,150	94.6 %
Pacific Ocean	7	19,835	4,867	-75.5 %	21,649	4,865	-77.5 %
Atlantic Ocean	12	14,301	3,378	-76.4 %	16,269	3,173	-80.5 %
Global Total	25	64,566	10,138	-84.3 %	77,435	10,188	-86.8 %

**IND-Table 1.** Indian Ocean localities of importance to *Eretmochelys imbricata* (n= 30), including 6 Index Sites (and their assigned reference numbers), for which quantitative data exist on past and present abundance (IND-Table 2). Long-term changes in population size were calculated with these data, and are presented in IND-Tables 3 and 4. IND-Table 5 presents current status and qualitative data pertaining to population trends for all 30 sites. Locations of the 6 Index Sites are shown in the map in W-Figure 1.

Index #	Nesting Sites	IND-Table(s)	Justification
<b>INDIAN OCEAN: SOUTH WESTERN</b>			
	Comoro Islands	5	
	France (Iles Esparse)	5	
1	Kenya	2, 3, 5	Surveys conducted in early 1980s and early 2000s
2	Madagascar	2, 3, 5	Well documented shell trade
	Mauritius	5	
	Mayotte	5	
	Mozambique	5	
3	Seychelles	2, 3, 4, 5	Nesting surveys conducted in early 1980s and early 2000s
	Tanzania	5	
<b>INDIAN OCEAN: NORTH WESTERN</b>			
	Bahrain	5	
4	Egypt	2, 3, 5	Beach surveys conducted in early 1980s and early 2000s
	Eritrea	5	
	Iran	5	
	Kuwait	5	
	Oman	5	
	Qatar	5	
	Saudi Arabia (Arabian Gulf)	5	
	Saudi Arabia (Red Sea)	5	
	Somalia	5	
	Sudan	5	
	United Arab Emirates	5	
	Yemen	5	

Index #	Nesting Sites	IND-Table(s)	Justification
<b>INDIAN OCEAN: CENTRAL &amp; EASTERN</b>			
	Australia: North West Shelf	5	
	British Indian Ocean Territory	5	
	India (Andaman & Nicobar Islands)	5	
	Malaysia (Melaka)	5	
5	Maldives	2, 3, 5	Well documented historic trade in hawksbill products.
	Myanmar	5	
6	Sri Lanka	2, 3, 5	Historic trade & recent beach survey data
	Thailand (Andaman Sea)	5	

**IND-Table 2.** Quantitative evaluation of nesting activity and population trends in the Indian Ocean based on available *Past* and *Recent* estimates for *Eretmochelys imbricata* at 6 sites. Data codes include: NF, numbers of nesting females; NN, numbers of nests; and TSE, Tortoiseshell Export Statistics. A bracketed figure of 3-5 nests per female was used to convert from number of nests to numbers of females. All values are based on annual means unless otherwise stated.

Index #	Index Nesting Site	Data type	Past Estimate 1		Past Estimate 2		Recent Estimate		Citation (Past)	Citation (Recent)
			Years	Mean	Years	Mean	Years	Mean		
INDIAN OCEAN: SOUTH WESTERN										
1	Kenya	NF	Late 1970s	50 females / yr			Early 2000s	< 10 females / yr	Frazier 1982	Okemwa <i>et al.</i> 2004
2	Madagascar	TSE	1870s	4,000 kg / yr	1928	1,440 kg / yr	1950's	1,000 kg / yr	Decary 1950 (cited in Hughes 1973); Petit 1930 (cited in Hughes 1973); Hughes 1973; Groombridge and Luxmoore 1989	Hughes 1973; Groombridge and Luxmoore 1989
		NF					2001	1,000 females / yr		A. Cooke, <i>in litt.</i> to J. Mortimer 2001
3	Seychelles <u>All</u> 22 Inner Islands	NF	Early 1980s	820 females /yr			Early 2000s	625 females / yr	Mortimer 1984, 2004	Mortimer 2004, 2006; See IND-Table 4 for details.
INDIAN OCEAN: NORTH WESTERN										
4	Egypt	NF	Early 1980s	200-500 females / yr			Early 2000s	50-100 females / yr	Frazier and Salas 1984	J.D. Miller, <i>in litt.</i> to J. Mortimer, 13 Nov 2006

Index #	Index Nesting Site	Data type	Past Estimate 1		Past Estimate 2		Recent Estimate		Citation (Past)	Citation (Recent)
			Years	Mean	Years	Mean	Years	Mean		
INDIAN OCEAN: CENTRAL & EASTERN										
5	Maldives	NN					1988-1995	2,300 nests/yr	Groombridge and Luxmoore 1989; Frazier <i>et al.</i> 2000	Zahir and Hafiz 1997
		NF	1970s	2,730 females /yr	mid-1980s	~500 females / yr		460-767 females / yr		
6	Sri Lanka South Coast	NN	1840s	Dense nesting reported	1990s	<10 nests / yr	2000s	<10 nests / yr	Bennett 1843; Amarasooriya 1996; Deraniyagala 1939	Kapurusinghe 2000
		NF		Estimated many hundreds of turtles		2-4 females / yr		2-4 females / yr		

**IND-Table 3.** Summary of estimated population change over 3 generations for the 6 Index Sites in the Indian Ocean. Figures derive from the Past and Recent Estimates presented in IND-Table 2, and from Exponential and Linear extrapolation functions (IUCN 2001a). Extrapolation functions are used only when there is a suspected change in the subpopulation size over a specific time interval outside of the period represented by data in IND-Table 2. Where bracketed estimates are presented in IND-Table 2, the mid-point is used here. In such cases, unless otherwise noted, both linear (L) and exponential (E) functions are used due to a lack of information on the true rate of change over the time interval. All values are based on annual means.

Index #	Index Sites	Raw Data (from IND-Table 2)		Notes on Population Trajectories & Comments on Current Status	Past Annual Nesting Female Subpopulation Size (3 generations back) (1870)	Present Annual Nesting Female Subpopulation Size (2005)	% Change over 3 generati ons
		Past	Present				
INDIAN OCEAN: SOUTH WESTERN							
1	Kenya	50 females/yr (1975)	10 females/yr (2005)	1870-1950: Exploited and probably declining 1950-2005: Declining	274 females/yr (L) (1870)	10 females/yr (R) (2005)	-96%
					378,201 females/yr (E) not realistic (1870)	10 females/yr (R) (2005)	na

Index #	Index Sites	Raw Data (from IND-Table 2)		Notes on Population Trajectories & Comments on Current Status	Past Annual Nesting Female Subpopulation Size (3 generations back) (1870)	Present Annual Nesting Female Subpopulation Size (2005)	% Change over 3 generati ons
		Past	Present				
2	Madagascar	Proxy:		1870-2005: Declining (Petit 1930, cited in Hughes 1973), Hughes 1973, Rakotonirina and Cooke 1994, A. Cooke, <i>in litt.</i> to J. Mortimer 2001). Some 4,000 kg shell exported annually from mid-1800s to 1920s; 1,440 kg in 1928; 1,000 kg in 1950s (Groombridge and Luxmoore 1989, Hughes 1973).  <i>Linear forward extrapolation of proxy data have resulted in unrealistic estimate for 2005. So, only exponential extrapolations of proxy data is used.</i>	Proxy:	-1,310 kg (L) <i>not realistic</i> (2005)	na
		4,000 kg/yr (1870s)	1,000 kg/yr (1950s)		3,935 kg (L) (1870)		
					3,697 kg (E) (1870)	353 kg (E) (2005)	- 90.5%
		Population estimate:			Extrapolated population estimate:		
		na	1,000 females/yr (2005)		na (L)	na (L)	na
					10,471 females/yr (E) (1870)	1,000 females/yr (R) (2005)	- 90.5%
3	Seychelles (All 22 Inner Islands)			1870-1970: Declining due to intensive shell trade (Mortimer 1984) 1971-2005: Declining at some islands and increasing at others, depending on management regime: 2 islands well protected since early 1970s; 7 islands with intermediate protection since 1979; and 13 islands with no protection prior to 1994 (Mortimer 2004, 2006). Since 1994, all turtles legally protected, but some poaching continues and unregulated coastal development threatens nesting habitat (Mortimer 2004).	1,922 females/yr (L) (1870)	605 females/yr (R) (2005)	-32%
		820 females/yr (1983)	625 females/yr (2003)				
					11,009 females/yr (E) (1870)	575 females/yr (R) (2005)	-95%



Index #	Index Sites	Raw Data (from IND-Table 2)		Notes on Population Trajectories & Comments on Current Status	Past Annual Nesting Female Subpopulation Size (3 generations back) (1870)	Present Annual Nesting Female Subpopulation Size (2005)	% Change over 3 generati ons
		Past	Present				
INDIAN OCEAN: NORTH WESTERN							
4	Egypt	350 females/yr (1980)	75 females/yr (2000)	1870-2005: Declining.	1,863 females/yr (L) (1870)	6 females/yr (L) (2005)	-99.7%
					1,754,239 females /yr (E) not realistic (1870)	50 females/yr (E) (2005)	na
INDIAN OCEAN: EASTERN							
5	Maldives	2,730 females/yr (1975)	614 females/yr (1992)	1870-1972: Long history of shell trade. Ban on eating hawksbill meat lifted in 1950 1973-1995: Ongoing exploitation for meat, shell and eggs 1995-2005: Moratorium on killing turtles implemented; but egg collection continues.	15,800 females/yr (L) (1870)	-1,003 females/yr (L) not realistic (2005) <614 females /yr (R) (2005)	-96%
				Exponential extrapolations produced unrealistic backward results. Linear regression produced unrealistic forward extrapolation.	24,768,300 females/yr (E) not realistic (1870)	176 females/yr (E) (2005)	na

Index #	Index Sites	Raw Data (from IND-Table 2)		Notes on Population Trajectories & Comments on Current Status	Past Annual Nesting Female Subpopulation Size (3 generations back) (1870)	Present Annual Nesting Female Subpopulation Size (2005)	% Change over 3 generati ons
		Past	Present				
INDIAN OCEAN: SOUTH WESTERN							
6	Sri Lanka South Coast	100s- 1,000s females/yr (1840s)	<2-4 females/yr (2000s)	1870-2005: Populations in decline due to heavy exploitation for shell, meat and eggs.	Minimum of 100 females/yr (R) (1870)	3 females/yr (R) (2005)	-96%
INDIAN OCEAN INDEX BEACHES:					3 Generations Back	Recent	% Change
TOTAL CHANGE USING RAW DATA + LINEAR FUNCTIONS					30,430	1,893	-93.8 %
TOTAL CHANGE USING RAW DATA + EXPONENTIAL FUNCTIONS (a)					39,517	2,150	-94.6 %

(a) For these exponential & raw data calculations, unreasonable exponential extrapolations produced for Kenya, Madagascar, Egypt, and Maldives were replaced by the more conservation linear extrapolations. It is noteworthy that when population declines over relatively short periods of time were steep, the exponential functions produced unrealistic extrapolated population sizes.

**IND-Table 4.** Changes in the size of hawksbill populations in Seychelles during two decades (from the early 1980s to the early 2000s) at islands that have had different management regimes since the 1970s (Source: Mortimer, 2004, 2006; Unpubl. data: Bird Island Lodge, Cousine Island Company, Denis Island, Fregate Island Private (FIP), International Council for Bird Preservation (ICBP), Island Conservation Society (ICS), J.A. Mortimer, Marine Conservation Society Seychelles (MCSS), Marine Parks Authority (SCMRT-MPA), Nature Seychelles, Nature Protection Trust of Seychelles (NPTS), North Island Seychelles, Royal Society for Nature Conservation (RSNC), and Seychelles Ministry of Environment & Natural Resources (MENR).

Management Regime of Nesting Populations	Data Type	Past Estimate		Recent Estimate		Changes in Population Size Over Two Decades
		Years	Mean	Years	Mean	
Seychelles <b>Well protected</b> since early 1970s (2 islands)	NF	Early 1980s	44 females /yr	Early 2000s	215 females /yr	+ 389 %
Seychelles <b>Intermediate protection</b> since 1979 (7 islands)	NF	Early 1980s	240 females /yr	Early 2000s	190 females /yr	- 21%
Seychelles <b>No protection</b> before 1994 (13 islands)	NF	Early 1980s	536 females /yr	Early 2000s	220 females /yr	- 59%
<b>Seychelles <u>All</u> 22 Inner Islands</b>	NF	<b>Early 1980s</b>	<b>820 females / yr</b>	<b>Early 2000s</b>	<b>625 females /yr</b>	<b>-24%</b>

**IND-Table 5.** Current population estimates and qualitative information about status and trends for reviewed hawksbill populations in the Indian Ocean. Population estimates are based on nesting females / yr, but where estimates are derived from numbers of nests, a bracketed figure of 3-5 nests per female is used to convert from numbers of nests to numbers of females.

Index #	Locality	Current Population Size	Comments (Source)	Status / Trends (Source)
<b>INDIAN OCEAN: SOUTH WESTERN</b>				
	<b>Comoro Islands</b>	25-50 females/yr	Ben Mohadji <i>et al.</i> 1996. Shell exports to Japan 1950-1990: 8,596 turtles (6,361 kg)	Population probably declining. On Grand Comore and Anjouan islands nesting habitat has been destroyed by sand mining (Ben Mohadji <i>et al.</i> 1996).
	<b>France</b> Iles Eparses  (Europa, Tromelin, Juan de Nova, Glorieuses)	20-45 females/yr	Europa: no nesting reported, but, immature foraging turtles occur (Gravier-Bonnet <i>et al.</i> 2006). Tromelin: estimated to have a few nesters Juan de Nova: estimated 10-30 females/yr Glorieuses: estimated <10 females/yr Source: J. Bourjea & S. Ciccione, <i>in litt.</i> to J. Mortimer, 10 Oct 2006.	Trends unknown.
1	<b>Kenya</b>	< 10 females/yr	Very sparse nesting, but significant foraging aggregations in Kenyan waters (Wamukoya <i>et al.</i> 1996, Okemwa <i>et al.</i> 2004). Shell exports 1970-1986 equivalent to 30,305 turtles.  Shell exports to Japan since 1950: 30,664 turtles (22,691 kg).	Remnant population. Declining. Unregulated coastal development threatens nesting habitat, and accidental mortality in fishing gear (esp. trawl nets and gill nets) a major threat to foraging aggregations (Okemwa <i>et al.</i> 2004). Trawler bycatch ~ 500-1,000 annually (Wamukoya <i>et al.</i> 1995) with Hawksbills being 6% of strandings in 2000-01 (Okemwa <i>et al.</i> 2004).

Index #	Locality	Current Population Size	Comments (Source)	Status / Trends (Source)
2	<b>Madagascar</b>	~1,000 females/yr	<p>Up to 1,000 females may nest annually, primarily on NE, NW and SW coasts (A. Cooke, <i>in litt.</i> to J. Mortimer 2001). Hughes (1973) estimated more than 2,500 Hawksbills killed annually, including ~600 adults. Surveys of &gt;20 islands in SW in 2001 show intense exploitation of nesting and foraging turtles and eggs (A. Cooke, <i>in litt.</i> to J. Mortimer 2001). Surveys of Nosy Hara-Radama Islands in NW in 2001 found nesting at uninhabited beaches, but frequent signs of opportunistic slaughter, relatively abundant foraging turtles, but net capture common (Metcalf <i>et al.</i> 2007). Four seasons of beach surveys (2000-04) at Nosy Iranja Kely in NW indicate sparse nesting (~20 nests/season/3 km beach) (Bourjea <i>et al.</i> <i>in press</i>).</p> <p>Shell exports to Japan since 1950: 1,808 turtles (1,338 kg).</p>	<p>Declining.</p> <p>Annual shell exports from mid-19<sup>th</sup> century through 1920 (Groombridge and Luxmoore 1989) were equivalent to 4,054-5,405 turtles. Drastic population declines after WWI reported (Petit 1930 as cited in Hughes 1973), and exports of 1,000 kg/yr (1,351 turtles) by mid-20<sup>th</sup> century, and 200 kg/yr (270 turtles) by 1973 (Hughes 1973). Sale of worked shell to tourists continues (Meylan and Donnelly 1999). Nesting turtles in surveyed areas appear to be in decline with exploitation for meat, eggs, and shell (Rakotonirina and Cooke 1994). Trawling along NW and W believed a threat (Randrianmiarana <i>et al.</i> 1998).</p>
	<b>Mauritius</b> (including St. Brandon)	< 50 females/yr	<p>Once abundant, now nest only at remote St. Brandon group (Frazier 1980); last recorded nesting on Mauritius in 1970s (Mangar and Chapman 1996). In 1996, all turtles encountered were killed; stuffed and tortoiseshell curios for sale in markets (Mangar and Chapman 1996). Turtles legally protected since 1998.</p>	Depleted.

Index #	Locality	Current Population Size	Comments (Source)	Status / Trends (Source)
	<b>Mayotte</b>	10-50 females/yr	<p>Nesting populations not yet adequately surveyed (Groombridge and Luxmoore 1989, M. Quillard &amp; S. Ciccione, <i>in litt.</i> to J. Mortimer 2006, J. Bourjea, <i>in litt.</i> to J. Mortimer 2006).</p> <p>Significant numbers foraging Hawksbills (Groombridge and Luxmoore 1989, M. Quillard &amp; S. Ciccione, <i>in litt.</i> to J. Mortimer 2006).</p>	<p>Trends unknown but believed to be declining. In early 1970s, turtles were killed whenever encountered (Frazier 1980). During past decade, poaching has continued, but public awareness campaigns underway (Cousin 2001, Quillard 2001).</p>
	<b>Mozambique</b>	<10 females/yr	<p>Nest in northern Mozambique, especially on offshore islands. During 1980s, eggs and meat taken extensively for subsistence. Currently very rare (A. Costa, <i>in litt.</i> to J. Mortimer 2006, J. Garnier <i>in litt.</i> to J. Mortimer 28 Mar 2007, I. Silva, <i>in litt.</i> to J. Mortimer 17 Apr 2007).</p> <p>Shell exports to Zanzibar recorded as early as 1890. Regular trade to Zanzibar during 1920-1964 and to France and Japan during 1965 to 1977 was approximately 20,700 kg (Frazier 1980), equivalent to 27,973 turtles.</p> <p>Shell exports to Japan since 1950: 985 turtles (729 kg).</p>	<p>Depleted and declining. Nest monitoring program at Vamizi and Rongui Islands suggest decline in recent years: 24 nests in 2003, 6 in 2004, 7 in 2005, 3 in 2006 (Silva and Garnier 2007, cited in J. Garnier, <i>in litt.</i> to J. Mortimer 28 Mar 2007).</p> <p>Recorded in fishing nets (A. Costa, <i>in litt.</i> to J. Mortimer 2006).</p>

Index #	Locality	Current Population Size	Comments (Source)	Status / Trends (Source)
3	<b>Seychelles</b> 22 Inner Islands	~ 625 females/yr	Shell export intensified in 19 <sup>th</sup> & 20 <sup>th</sup> centuries (Mortimer 1984). In mid-1960s through mid-1990s most females killed before reproducing at unprotected beaches (Mortimer 1984, 1998). Entanglement in gill nets most significant fisheries related threat (Domingue and Mortimer 2001). Since 1994, sea turtles legally protected (Mortimer 1998) and domestic tortoiseshell trade ceased (Mortimer 1999). Shell exports from Seychelles to Japan since 1950: 8,877 turtles (6,569 kg).	Depleted and declining at unprotected sites. Nesting population depleted by early 1980s (Mortimer 1984); overall population declines continued through early 2000s. Declines at unprotected and poorly protected sites; increases at several well protected sites (Mortimer 2004, 2006) (see IND-Table 4). Inadequate control of coastal development seriously threatens nesting habitat (Mortimer 2004).
	<b>Seychelles</b> Outer Islands	~ 800 females/yr	Estimate based on national surveys conducted in early 1980s (Mortimer 1984) and recent unpublished data (Mortimer unpubl. data).	Depleted. Overall trend unknown. Unpublished data suggest increases at protected sites and declines at unprotected islands.

Index #	Locality	Current Population Size	Comments (Source)	Status / Trends (Source)
	<b>Tanzania</b>	<50 females/yr	<p>Foraging animals more abundant than nesters. Zanzibar is historically a major clearing house. From 1891 to 1963, ~325 kg were exported annually from the mainland to Zanzibar (Frazier 1980). For the period 1891 to 1950, these exports represent 26,351 turtles. Tanzania, including Zanzibar, was the largest supplier of shell to Japan in the western Indian Ocean from 1950-1986 (Milliken and Tokunaga 1987, Groombridge and Luxmoore 1989)</p> <p>In 1970s estimated 50 nesters/yr, mostly at Mziwi Island (Frazier 1982) which has since sunk (Howell and Mbindo 1996). Turtles caught on feeding grounds, while nesting, and by dynamite fishing (Frazier 1982).</p> <p>Shell exports to Japan since 1950: 65,001 turtles (48,101 kg).</p>	<p>Depleted and declining.</p> <p>Threats include unregulated coastal development and incidental capture in fishing gear (Frazier 1980, Howell and Mbindo 1996).</p>
<b>INDIAN OCEAN: NORTH WESTERN</b>				
	<b>Bahrain</b>	Sparse	Data lacking.	Trends unknown.
4	<b>Egypt</b>	~50-100 females/yr	<p>Recent estimate of 50-100 females annually (J. Miller <i>in litt.</i> to J. Mortimer, 13 Nov 2006), is lower than the 200-500 reported by Frazier and Salas (1984). Historically important source and consumer of shell (Parsons 1972). Most of nesting on offshore islands (Frazier and Salas 1984).</p>	<p>Declining.</p> <p>Destruction of habitat from oil pollution, underwater explosions related to seismic oil exploration problematic in 1980s (Frazier and Salas 1984). Current threats include coastal development and near shore reef habitat destruction (J.D. Miller, <i>in litt.</i> to J. Mortimer, 13 Nov 2006).</p>



Index #	Locality	Current Population Size	Comments (Source)	Status / Trends (Source)
	<b>Eritrea</b>	unknown	<p>No estimate available (Hillman and Gebremariam 1996).</p> <p>Shell exports to Japan since 1950: 4,809 turtles (3,559 kg)</p>	<p>Status and trends unknown.</p> <p>Sparse coastal human population indicates neither subsistence take nor coastal development likely to pose a threat (Hillman and Gebremariam 1996). Fisheries related mortality (esp. trawlers and shark nets) may be a serious problem with an estimated 0.61 turtles (47% are Hawksbills) caught per hour trawled in Eritrean waters (Gebremariam <i>et al.</i> 1998).</p>
	<b>Iran</b>	~500-1000 females/yr	<p>Historic data from the 1970s indicate Shidvar, Lavan, Hormuz, Larak, Queshm, and Jabrin islands and adjacent mainland beaches hosted significant, but poorly surveyed nesting populations (Kinunen and Walczak 1971, Ross and Barwani 1982). Recent data indicate that Ommolkaram and Nakhiloo islands of Booshehr Province (Valavi 1999; J. Mortimer, pers. obs, 2001; Mobaraki 2003, 2004a; Mobaraki and Elmi 2005), Shidvar and Hendourabi islands of Hormozgan Province, and Nayand Bay, (Mobaraki 2003, 2004a, 2004b) are important sites.</p>	<p>Trends unknown.</p> <p>Populations threatened by egg collection and predation, especially on mainland (Mobaraki 2004); killing of nesting females (Mobaraki 2004a), incidental capture in fishing gear (J. Mortimer, pers. obs. 2001, Mobaraki and Elmi 2005). Foraging habitat degradation due to coral bleaching events (Sheppard and Loughland 2002, Sheppard 2006) and oil spills (Miller 1989).</p>
	<b>Kuwait</b>	<20 females/yr	<p>Small amount of nesting occurs on Um Al-Maradm and Garu islands (Groombridge and Luxmoore 1989).</p>	<p>Trends unknown.</p>

Index #	Locality	Current Population Size	Comments (Source)	Status / Trends (Source)
	<b>Oman</b>	~ 600-800 females/yr	<p>Nesting primarily on coast of Gulf of Oman (Salm <i>et al.</i> 1993, Baldwin and Al-Kiyumi 1997) including: 250-350 at the protected Dimaniyat Islands; and 100 at Masirah island (Ross and Barwani 1982, Ross 1981).</p> <p>Salm <i>et al.</i> (1993) considered Dimaniyat the most important Hawksbill sanctuary in region.</p>	<p>Possibly stable.</p> <p>Monitoring at Dimaniyat Islands indicates stable nesting numbers (pers. comm. A. Al-Kiyumi to N. Pilcher 2006). Egg collection reported at Masirah, Bar al Hikman, and Dimaniyat Islands (Salm 1991 cited in Baldwin and Al Kiyumi 1997). On mainland beaches foxes destroyed 62-82% of eggs, and 10-15% were laid below high tide line (Salm 1991 cited in Baldwin and Al Kiyumi 1997). Currently, main threats are incidental capture in nets, loss of nesting habitat &amp; disturbance on nesting beaches (R. Baldwin in litt. to E. Possardt, 20 Jun 2007). Other problems include rainwater runoff, tourist activities, and vehicular traffic (Baldwin and Al-Kiyumi 1997, Rees and Papathanasopoulou 2006).</p>
	<b>Qatar</b>	estimated >100 females/yr	<p>Nesting reported Ras Laffan (Pilcher 2006), and Sharaawh and Dayinah islands (Ross and Barwani 1982). Development of Port at Ras abu Khamis destroyed coral reefs and Hawksbill population (Ross and Barwani 1982). During 1970s, meat and eggs commonly eaten (Frazier 1980).</p>	<p>Stable.</p> <p>During six years (2001-06) of monitoring, the Ras Laffan population appears to be stable with an average of 178 nests/yr (Pilcher 2006). Nesting populations threatened by habitat degradation that includes dead corals killed in 1998 and 2000 bleaching events (N. Pilcher, <i>in litt.</i> to J. Mortimer, 13 Aug 2006), oil pollution and lighting issues.</p>

Index #	Locality	Current Population Size	Comments (Source)	Status / Trends (Source)
	<b>Saudi Arabia</b> Arabian Gulf	~ 175-265 females/yr	<p>Estimates by island based on Pilcher (1999) and J.D. Miller (<i>in litt.</i> to J. Mortimer, 13 Nov 2006: Jana, 100-150; Karan, &lt;50; Jurayd, 10-15; and Kurayn, &lt;50.</p> <p>Shell exports from Saudi Arabia to Japan since 1950: 149 turtles (110 kg).</p>	<p>Trends unknown.</p> <p>Saudi nationals do not eat turtle eggs or meat, but foreigners on fishing boats do (Pilcher 1999). Gill nets entangle hatchlings on beach (Pilcher 1999) and turtles in water (Miller 1989). The most serious threat is destruction of nesting and foraging habitats. Tar, oil slicks and debris on shore entrap hatchlings and prevent nesting (Miller 1989, Pilcher 1999). Spilled oil and dispersants threaten marine ecosystems (Miller 1989). Coral bleaching events in 1998 and 2000 destroyed much coral reef in The Gulf (Sheppard and Loughland 2002, Sheppard 2006). Pilcher (1999) cites need for regular patrol of nesting beaches to address threats.</p>
	<b>Saudi Arabia</b> Red Sea	100-200 females/yr	<p>Estimate based on J.D. Miller (<i>in litt.</i> to J. Mortimer, 13 Nov 2006). Low density nesting occurs at numerous sites from the islands of the Farasan Archipelago to Tiran Island at the Gulf of Aqaba (Miller 1989).</p>	<p>Trends unknown.</p> <p>Major threats identified by Miller (1989; <i>in litt.</i> to J. Mortimer, 13 Nov 2006) include: egg collection; fisheries related mortality (esp. trawlers), and habitat destruction caused by cement dust.</p>
	<b>Somalia</b>	unknown	<p>Nesting reported in NE zone and SW regions, but no estimates available (Abdulrizak Osman Ali, Ocean Training Promotion, pers. comm. to J. Mortimer, 2000). Bajun on south coast exploited shell for generations; sold to Europe in 1970s, and formerly to Zanzibar at ~100 kg/yr, except for 5,099 kg exported in 1976 (Frazier 1980).</p> <p>Shell exports to Japan since 1950: 2,407 turtles (1781 kg).</p>	<p>Trends unknown.</p>

Index #	Locality	Current Population Size	Comments (Source)	Status / Trends (Source)
	<b>Sudan</b>	300-350 females/yr	<p>Estimate based on 1970s data. Most nesting restricted to distant islands in Suakin Archipelago (Moore and Balzarotti 1977, Hirth and Abdel Latif 1980) and islands off Mohammed Qol (Moore and Balzarotti 1977).</p> <p>Formerly intense tortoiseshell trade (Groombridge and Luxmoore 1989). Killed in large numbers for meat in late 19<sup>th</sup> century at opening of Suez Canal (Moore and Balzarotti 1977). Subsistence take in 1970s (Frazier 1980).</p>	Depleted. Trends unknown.
	<b>United Arab Emirates</b>	100-200 females/yr	Estimate based on J.D. Miller ( <i>in litt.</i> to J. Mortimer, 13 Nov 2006). Nesting occurs at offshore islands	<p>Trends unknown.</p> <p>Current threats include incidental capture in fish traps and set-nets (J.D. Miller, <i>in litt.</i> to J. Mortimer, 13 Nov 2006).</p>
	<b>Yemen</b>	~500 females/yr ??	<p>Estimate based on data from 1960s and 1970s. Nesting reported for Socotra, Abd al Kuri, Jabal Aziz and Perim, and at low coral islands 3-30 km offshore (Hirth 1968 as cited in Ross and Barwani 1982, Groombridge and Luxmoore 1989). Meat and eggs eaten by fishermen (Frazier 1980).</p> <p>Shell exports to Japan since 1950: 49 turtles (36 kg).</p>	Trends unknown.

Index #	Locality	Current Population Size	Comments (Source)	Status / Trends (Source)
<b>INDIAN OCEAN: CENTRAL &amp; EASTERN</b>				
	<b>Australia</b> Western Australia (WA)	~ 2,000 females/yr (rough estimate)	WA Hawksbill population is the largest in the Indian Ocean (Limpus 1997, 2002), represented by genetic stock centered on Rosemary Island in Dampier Archipelago, site of long term tagging project (Broderick <i>et al.</i> 1994, Prince 1994 cited in Pendoley 2005). Nesting distribution has been mapped, but population sizes poorly quantified (Limpus 2002). Montebello Group now an important rookery for hawksbills, was the site of three nuclear tests in the 1950s, resulting in 'tens of thousands' of dead and rotting turtles (no species ID) on the beach (Pendoley 2005). Shell exports from Australia to Japan since 1950: 29,109 turtles (25,616 kg).	Status and trends unknown. Much WA nesting occurs within areas of greatest industrial development, including brightly lit oil/gas facilities on islands and at sea (Pendoley 2005, Limpus 2002). Altered light horizons may reduce nesting activity and increase hatchling predation at sea. No monitoring of expanding human populations, new holiday huts on nesting islands, and associated habitat destruction, increased boat strikes, and other disturbances (Limpus 2002). Satellite tracking of post nesting WA Hawksbills indicated migrations of 50-450 km from nesting beach into unprotected waters (Pendoley 2005), suggesting possibility of mortality similar to that of eastern Australian populations (Limpus 2002).
	<b>British Indian Ocean Territory</b> (Chagos Islands)	~ 300-700 females/yr	Inhabited from 1780s until 1972 when US/UK military base was established at Diego Garcia. Historical records show "significant export" of tortoiseshell (Parsons 1972), but during 20 <sup>th</sup> century (1904-1929) annual take was less than 200 animals /yr (Frazier 1980). Diego Garcia hosts the most important nesting and foraging habitats, and offers good protection (Mortimer and Day 1999, Mortimer 2000), but poaching continues in the outer islands (Mortimer, unpublished data, 2006).	Depleted. Current trend unknown. Significant decline since late 18 <sup>th</sup> century. Turtles now protected by law (Mortimer and Day 1999), but enforcement difficult in outer islands (Mortimer unpubl. data, 2006). Current population trend unknown. Erosion of nesting beaches is serious long-term problem, especially in outer islands (Mortimer and Day 1999, Mortimer unpubl. data, 2006), perhaps due to sea level rise and coral reef mortality (Sheppard 2002).

Index #	Locality	Current Population Size	Comments (Source)	Status / Trends (Source)
	<b>India</b> (Andaman & Nicobar Islands)	~250 females/yr	Incomplete surveys conducted in 1992 estimate 205 females nesting annually in Andaman Islands at 30 sites, and 45 females in Nicobar group at 11 sites (Andrews <i>et al.</i> 2006). Shell exports to Japan since 1950: 7,868 turtles (5,822 kg).	Declining. Threats include sand mining, egg predation by dogs and pigs, incidental capture in active and discarded gill nets, and poaching of nesting females and foraging turtles by settlers (Andrews <i>et al.</i> 2006).
	<b>Malaysia</b> Melaka	~ 50-85 females/yr	Prior to 1990 when Department of Fisheries Malaysia established hatchery, people consumed most eggs (Mortimer <i>et al.</i> 1993). Melaka coastline now undergoing intensive coastal development and massive land reclamation (Min Min Lau, pers. comm. to J. Mortimer; J. Mortimer, pers. obs., 2003). Shell exports to Japan from West Malaysia since 1950: 21,169 turtles (15,665 kg). (Records do not specify from which State(s) in West Malaysia the shell originated).	Depleted. Numbers of eggs incubated per year during 1991-2005 have remained stable, averaging ~250 egg clutches/ yr (source: Department of Fisheries Malaysia); but, apparent stability may reflect increased efforts to protect despite possible decline (Min Min Lau (WWF-M), pers. comm. to J. Mortimer 2006). Current threats include destruction of nesting habitat and entanglement in fishing nets (Min Min Lau, pers. comm. to J. Mortimer, 2006). Recently Malaysian Fisheries Department purchased the island of Pulau Upeh, the most important nesting site in Melaka (Mortimer <i>et al.</i> 1993), to make it a turtle sanctuary (Lee 2006).

Index #	Locality	Current Population Size	Comments (Source)	Status / Trends (Source)
5	<b>Maldives</b>	~ 460-767 females/yr	<p>Estimate based on data collected in 1980s (Frazier <i>et al.</i> 2000) and during 1988-95 (Zahir and Hafiz 1997). 20<sup>th</sup> Century exploitation intense for shell, meat and eggs. 65% of Hawksbills in trade were caught on beach (Groombridge and Luxmoore 1989). In 1995, a 10 year moratorium on killing turtles was implemented but eggs were not protected. In 2006, egg protection was enacted at 11 islands and will be expanded to 13 islands in 2007 (out of several thousand islands in country) (H. Zahir, <i>in litt.</i> to J. Mortimer, 19 Jan 2006).</p> <p>Shell exports to Japan 1950-1990: 28,141 turtles (20,824 kg) but major exports were from 1985 onward.</p>	<p>Declining.</p> <p>Long history of tortoiseshell export combined with hunting for eggs and meat had tremendous impact (Frazier 1980). In early 1980s Maldives considered one of most important areas for Hawksbills in Indian Ocean, but exploitation identified as probable cause for depletion (Groombridge 1982). Continued decline likely because: a) No protected nesting areas in Maldives; b) No regulation of egg collection until 2006 when 11 islands protected (H. Zahir, <i>in litt.</i> to J. Mortimer, 19 Jan 2006).</p>
	<b>Myanmar</b>	less than 5 females/yr	<p>Estimate based on data from Maxwell (1911 as cited in Groombridge and Luxmoore 1989), ~100 nests/yr on one island off the Bawmi Circle in the Bassein District.</p>	<p>Remnant population. Declining.</p> <p>Probably shared decline of Burmese sea turtles of 90% over past century due to egg over-exploitation (Groombridge and Luxmoore 1989).</p>

Index #	Locality	Current Population Size	Comments (Source)	Status / Trends (Source)
6	<b>Sri Lanka</b> South coast	~ 10 females/yr	<p>Current estimate from T. Kapurusinghe (<i>in litt.</i> to J.A. Mortimer, 2006). Nesting along South coast so abundant in mid-19<sup>th</sup> century that Government sold individuals the right to capture them; and a flourishing local artisanal trade developed (Deraniyagala 1939). Legislation protecting turtles and eggs enacted in 1972, but ignored (Hewavisenthi 1990, Salm 1981 as cited in Groombridge and Luxmoore 1989). Juveniles still occur in offshore waters. Heavy exploitation continues. Virtually no egg survival outside hatcheries, except for in situ protection conducted by TCP (Kapurusinghe 2000). Many hatcheries poorly managed.</p> <p>Shell exports to Japan since 1950: 85 turtles (63 kg).</p>	<p>Remnant population. Declining.</p> <p>Historically important center for the tortoiseshell trade. Nesting Hawksbills abundant in 1840s (Deraniyagala 1939). No significant nesting remains; only six nests recorded by TCP during 1996-2000 from Rekawa beach (Kapurusinghe, 2000). From December 2006 to March 2007, TCP protected 20 Hawksbill nests; additional nests may be in unsurveyed areas of Sri Lanka (L. Ekanayake, <i>in litt.</i> to J. Mortimer, 2007).</p>
	<b>Thailand</b> Andaman Sea coast	< 10 females/yr	<p>Exploitation for meat &amp; eggs unregulated until 1947 Fisheries Act which prohibited killing of turtles &amp; established egg concession system requiring protection of 10-15% of eggs (Mortimer 1988). By 1980s, egg concessions abandoned at most sites due to disturbance from massive coastal development and tourism, but several National Parks established (Mortimer 1988). Major threats include poaching of eggs and turtles by Moken ("sea gypsy") people, and fisheries related mortality (Mortimer 1988; M. Aureggi, <i>in litt.</i> to J. Mortimer, 21 Aug 2006).</p> <p>Shell exports to Japan since 1950: 27 turtles (20 kg).</p>	<p>Remnant population. Declining.</p> <p>In 1980s small numbers of Hawksbills nested at Sulin and Similan islands, Phang Nga Province, and at Tarutao National Park in Satun Province (Ginsberg and Congdon 1981, cited in Mortimer 1988, Mortimer 1988). Hawksbill nesting is now reported only at Ko Surin National Park, where eggs collected by sea gypsies are sold to park officers for incubation--but with poor hatch success (M. Aureggi, <i>in litt.</i> to J. Mortimer, 21 Aug 2006).</p>



**PAC-Table 1.** Pacific Ocean localities of importance to *Eretmochelys imbricata* (n= 20), including 7 Index Sites (and their assigned reference numbers), for which quantitative data exist on past and present abundance (PAC-Tables 2 and 4). Long-term changes in population size were calculated with these data, and are presented in PAC-Tables 3 and 4. PAC-Table 5 presents current population status and qualitative data pertaining to population trends for all 20 sites. Locations of the 7 Index Sites are shown on the map in W-Figure 1.

Index #	Nesting Sites	PAC-Table(s)	Justification
PACIFIC OCEAN: WESTERN			
7	Australia: Torres Strait-Northern Great Barrier Reef (Milman Island = Index Beach)	2, 4, 5	Milman Island monitoring since 1990
	Australia: Northeastern Arnmem Land	5	
8	Indonesia	2, 3, 4, 5	Monitoring during both mid-1980s and during 1995-2005 at 14 sites
9	Japan	5	Monitoring underway since 1979
	Malaysia (East): Sabah Turtle Islands	2, 4, 5	
10	Malaysia (West): Terengganu	2, 4, 5	Monitoring underway since 1978
	Papua New Guinea	5	
	Philippines	5	
12	Thailand (Gulf of Thailand)	2, 4, 5	Data from mid-1950s; Monitoring underway from 1973-2005
	Vietnam	5	
PACIFIC OCEAN: CENTRAL			
11	American Samoa and Western Samoa	5	Monitoring during 1960s, and intermittently from 1992-2005
	Fiji	5	
	Guam	5	
	Hawaii	5	
	Micronesia	5	
	Palau Republic	5	
	Solomon Islands	2, 4, 5	
	Vanuatu	5	
	PACIFIC OCEAN: EASTERN		
13	El Salvador	5	Historical records and recent monitoring programs
	Mexico	2, 4, 5	

**PAC-Table 2.** Quantitative evaluation of nesting activity and population trends in the Pacific Ocean based on available *Past* and *Recent* estimates for *Eretmochelys imbricata* at 7 sites. Data codes include: NF, numbers of nesting females; NN, numbers of nests; FA number of foraging animals; and UPE, unit patrol effort at the nesting beach. A bracketed figure of 3-5 nests per female was used to convert from number of nests to numbers of females. All values are based on annual means unless otherwise stated.

Index #	Index Nesting Site	Data type	Past Estimate 1		Past Estimate 2		Recent Estimate		Citation (Past)	Citation (Recent)
			Years	Mean	Years	Mean	Years	Mean		
PACIFIC OCEAN: WESTERN										
7	Australia (Northern Territory & Queensland) Milman Island (Standard one month census used as Index for Torres Strait-Northern Great Barrier Reef sub-population)	NF	1990-1995	304 females/yr			1996-1999	292 females/yr	Miller <i>et al.</i> 2000	Miller <i>et al.</i> 2000
8	Indonesia <i>Summary of 14 Sites from PAC-Table 4)</i>	NN	1980s	8,113 nests/yr			1995 - 2006	2,630 nests/yr	See PAC-Table 4 for details	See PAC-Table 4 for details
		NF		1,623-2,704 females/yr				526-877 females/yr		
9	Malaysia SabahTurtle Islands Park	NN	1979-1987	381 nests/yr	1988-1996	443 nests/yr	1997-2005	347 nests/yr	Sabah Parks unpub. data; Groombridge and Luxmoore 1989, Pilcher and Ali 1999.	Sabah Parks unpub. data; P. Basinthal in litt. to J. Mortimer, 2006.
		NF		76-127 females/yr		89-48 females/yr		69-116 females/yr		

Index #	Index Nesting Site	Data type	Past Estimate 1		Past Estimate 2		Recent Estimate		Citation (Past)	Citation (Recent)
			Years	Mean	Years	Mean	Years	Mean		
10	Malaysia Terengganu State	NN	1978	69 nests/yr	1984-1991	41 nests/yr	1992-2000	18 nests/yr	Siow and Moll 1982, Chan and Liew 1999, Liew 2002	Liew 2002
		NF		14-23 females /yr		8-14 females/yr		4 -6 females/yr		
12	Thailand Ko Khram	NN	1956	224 nests in 1956	1973-1989	84 nests/yr	1990-2005	56 nests/yr	Monanunsap 1997, Groombridge and Luxmoore 1989, Charuchinda and Monanunsap 1998; Unpubl. data, M. Charuchinda; M. Aureggi, in litt.. to J. Mortimer, 13 Oct 2006	Charuchinda and Monanunsap 1998, Unpubl. data, M. Charuchinda, M. Aureggi, in litt.. to J. Mortimer, 13 Oct 2006
		NF	mid-1950s	45-75 females/yr		17-28 females/yr		11-19 females/yr		

Index #	Index Nesting Site	Data type	Past Estimate 1		Past Estimate 2		Recent Estimate		Citation (Past)	Citation (Recent)
			Years	Mean	Years	Mean	Years	Mean		
PACIFIC OCEAN: CENTRAL										
11	Solomon Islands Arnavon Community Marine Conservation Area (ACMCA)	UPE	1960s	Average of 14.0 nests /night at peak season ( ~100 nests per week at peak season; at Sikopo island 20 females taken in two nights.)	1992-1995; 2000	Average of 1.3 to 2.1 nests / night.	2005-2006	Average of 2.9 to 3.9 nests/night	McKeown 1977, Vaughn 1981, Ramohia and Pita 1996.	P. Ramohia and C. Siota, in litt. to J. Mortimer (28 Aug 2006; 3 Nov 2006)
		NN					2000-2005	500-600 nests/yr		Ramohia and Pita 1996, Mortimer 2002.
		NF						100-200 females/yr		
PACIFIC OCEAN: EASTERN										
13	Mexico Baja California	FA	1950s	foraging animals abundant	late 1960s	foraging animals rare or absent	1998 - 2001; 2005	foraging animals are rare or absent	Aschmann 1966 (cited in Seminoff <i>et al.</i> 2003b), Felger and Moser 1985 (cited in Seminoff <i>et al.</i> 2003b)	Seminoff <i>et al.</i> 2003b, J. Nichols, unpubl. data.

**PAC-Table 3.** Summary of estimated population change over 3 generations for the 7 Pacific Ocean Index Sites. Figures derive from the Past and Recent Estimates presented in PAC-Table 2, and from Exponential and Linear extrapolation functions (IUCN 2001a). Extrapolation functions are used only when there is a suspected change in the subpopulation size over a specific time interval outside of the period represented by data in PAC-Table 2. Where bracketed estimates are presented in PAC-Table 2, the mid-point is used here. In such cases, unless otherwise noted, both linear (L) and exponential (E) functions are used due to a lack of information on the true rate of change over the time interval. All values are based on annual means.

Index #	Subpopulation (Index Site)	Raw Data (from PAC-Table 2)		Notes on Population Trajectories & Comments on Current Status	Past Annual Nesting Female Subpopulation Size (3 generations back)	Present Annual Nesting Female Subpopulation Size (2005)	% Change over 3 generations
		Past	Present				
PACIFIC OCEAN: WESTERN							
7	Australia (Torres Strait-Northern Great Barrier Reef sub- population)  Milman Island (standard 1 month census data)	304 females/yr (1990-1995)	292 females/yr (1996-1999)	1870-1915: Trend unknown. 1916-1930s: Possibly declining due to intensive exploitation for shell (Limpus 2004). 1930s-1990: Shell industry ceased during 1930s; Hawksbill protected in Queensland in 1968. Trend unknown. 1991-2005: 3-4% annual population decline (Limpus and Miller 2000, Limpus pers. comm to J.A. Mortimer).  Projected decline of >90% from 1990 to 2020 (Limpus 2004)	Conservative backward extrapolation only to 1970:		
					445 females/yr (L) (1970)	239 females/yr (L) (2005)	-46%
					444 females/yr (E) (1970)	228 females/yr (E) (2005)	-49%
					Backward extrapolation for entire northern Australia subpopulation based on conservative backward extrapolation of Milman Island data (only to 1970):		
					7,448 females/yr (L) (1970)	4,000 females/ yr (L) (2005)	-46%
					7,789 females/yr (E) (1970)	4,000 females/ yr (E) (2005)	-49%

Index #	Subpopulation (Index Site)	Raw Data (from PAC-Table 2)		Notes on Population Trajectories & Comments on Current Status	Past Annual Nesting Female Subpopulation Size (3 generations back)	Present Annual Nesting Female Subpopulation Size (2005)	% Change over 3 generations
		Past	Present				
8	<b>Indonesia</b> <u>Summary of 14 Sites from PAC-Table 4)</u>	<b>2,164</b> females/yr (1985)	<b>701</b> females/yr (2005)	<p><b>1870-1917:</b> Declining due to long term shell trade (Parsons 1972).</p> <p><b>1918-1992:</b> Declining due to exploitation for eggs and shell (Dammerman 1929, Schulz 1987, Milliken and Tokunaga 1987).</p> <p><b>1993-2005:</b> Ongoing egg exploitation (Suganuma <i>et al.</i> 1999, H. Suganuma <i>in litt.</i> to J. Mortimer 2006)</p> <p><i>Exponential extrapolations produced unrealistic backward results.</i></p>	<p><b>10,571</b> females/yr (L) (1870)</p> <p><b>1,514,024</b> females/yr (E) <b>not realistic</b> (1870)</p>	<p><b>701</b> females/yr (R) (2005)</p> <p><b>701</b> females/yr (R) (2005)</p>	<p><b>-93%</b></p> <p><b>na</b></p>
9	<b>Malaysia</b> Sabah Turtle Islands Park	<b>102</b> females/yr (1985)	<b>92</b> females/yr (2005)	<p><b>1870-1946:</b> Population trend unknown.</p> <p><b>1947-1970:</b> Over exploitation of eggs and destruction of nesting beach from sand mining (de Silva 1982). Probable population decline.</p> <p><b>1971-1977:</b> Establishment of Marine Turtle National Park.</p> <p><b>1971-1984:</b> Population trends unknown.</p> <p><b>1985-2005:</b> Nesting numbers stable.</p>	<b>unknown</b> (1870)	<b>92</b> females/yr (R) (2005)	<p><b>Unknown trend</b> (decline likely since 1870; but apparently stable since 1985)</p>

Index #	Subpopulation (Index Site)	Raw Data (from PAC-Table 2)		Notes on Population Trajectories & Comments on Current Status	Past Annual Nesting Female Subpopulation Size (3 generations back)	Present Annual Nesting Female Subpopulation Size (2005)	% Change over 3 generations
		Past	Present				
10	Malaysia Terengganu State	18.4 females/yr (1978)	7.5 females/yr (2000)	1870-1950: Population trend unknown; 1951-2005: Population declining.	Conservative backward extrapolation only to 1950:  89.4 females/yr (L) (1870) (based on backward extrapolation to 1950)	3.8 females/yr (2005)	- 96%
					1,562 females/yr (E) (1870) (based on backward extrapolation to 1950)	1.1 females/yr (E) (2005)	-99%
12	Thailand Ko Khram	60 females/yr (1956)	15.1 females/yr (2001-2005)	1870-1900: Trend unknown. 1901-1947: Population declining. 1947: Establishment of Fisheries Regulations. 1948-2005: Continuing decline.	Conservative backward extrapolation only to 1901:  60 females/yr (L) (1870) (based on backward extrapolation to 1901)	12.3 females/yr (L) (2005)	- 80%
					60 females/yr (E) (1870) (based on backward extrapolation to 1901)	12.8 females/yr (E) (2005)	- 81%

Index #	Subpopulation (Index Site)	Raw Data (from PAC-Table 2)		Notes on Population Trajectories & Comments on Current Status	Past Annual Nesting Female Subpopulation Size (3 generations back)	Present Annual Nesting Female Subpopulation Size (2005)	% Change over 3 generations
		Past	Present				
PACIFIC OCEAN: CENTRAL							
11	Solomon Islands Arnavon Community Marine Conservation Area (ACMCA)	Proxy:		1870-1995: Population declining, due to shell trade. 1996: ACMCA established. 1997-2005: Decline arrested and possibly reversed.  <i>Exponential extrapolations produced unrealistic backward results.</i>	Proxy (# nests / night at peak season):		
		14.0 nests/night (1960s)	3.7 nests/night (2005)		39.4 nests/night (L) (1870)	3.7 nests/night (L) (2005)	-91%
					271.1 nests/night (E) (1870)	3.0 nests/night (E) (2005)	- 99%
		Population estimate:			Population estimate:		
		na	150 females/yr (2005)		1,667 females/yr (L) (1870)	150 females/yr (R)	-91%
					15,000 females/yr (E) <i>not realistic</i> (1870)	150 females/yr (R)	na
PACIFIC OCEAN: EASTERN							
13	Mexico Baja California	foraging animals abundant (1950s)	foraging animals rare or absent (2005)	1870-2005: Significant decline (Seminoff et al., 2003b)	abundant (1870)	rare or absent (2005)	- 80% ?
PACIFIC OCEAN INDEX BEACHES:					3 Generations Back	Recent	% Change
TOTAL CHANGE USING RAW DATA + LINEAR FUNCTIONS					19,835	4,867	-75.5 %
TOTAL CHANGE USING RAW DATA + EXPONENTIAL FUNCTIONS <sup>(a)</sup>					21,649	4,865	-77.5 %

(a) For these exponential + raw data calculations, unreasonable exponential extrapolations produced for Indonesia and Solomon Islands (AMCMA) were replaced by the more conservative linear extrapolations. It is noteworthy that when population declines over relatively short periods of time were steep, the exponential functions produced unrealistic extrapolated population sizes.



**PAC-Table 4.** Detailed quantitative evaluation of nesting activity and population trends at 18 sites in Indonesia in the Western Pacific Ocean based on available *Past* and *Recent* estimates for *Eretmochelys imbricata*. Data codes include: NN, numbers of nests. All values are based on annual means. Index Sites include those 14 sites for which both "Past Estimate 1" data collected during the 1980s, and "Recent Estimate" data collected between 1995 and 2005 are available. Past and Recent data from these 14 Index Sites are considered to be comparable, and "Population Trend" data are indicated by total figures for "Past Estimate 1" and "Recent Estimate" at the bottom of the table, and also feature in PAC-Tables 2 and 3.

For each site for which recent estimates are available (N=17), "Current Protected Status" is indicated along with the numbers of nests protected annually and a code indicating which organisation(s) implement protection: A = Everlasting Nature of Asia (ELNA); B = Indonesia Sea Turtle Research Center (ISTRIC); C = WWF-Indonesia; D = Directorate General of Protection and Nature Conservation (PHPA); E = Seribu National Park Rangers; F = Japan Bekko Association; G = Alas Purwo National Park.

Data from the four sites that have not been assigned Index # (and whose rows are shaded by grey) are not directly comparable to those of the 14 Index Sites, and so are not included in the totals at the bottom of this table summarising "Population Trends". But, for three of those four sites recent data are available, and these are included in the "Current Protected Status" summary at the bottom of the table.

Index #	Nesting site	Data type	Past estimate 1		Past estimate 2		Recent estimate		Currently protected?		Citation (past)	Citation (recent)
			Years	Mean	Years	Mean	Years	Mean	Yes/No	# Nests		
PACIFIC OCEAN: WESTERN												
Indonesia												
8-a	(Bangka-Belitung Province) Langkuas Islands	NN	1980s	100 nests/yr			1995-1997	< 50 nests/yr	No		Groombridge and Luxmoore 1989	Suganuma <i>et al.</i> 1999
8-b	(Bangka-Belitung Province) Lima Islands	NN	1980s	300 nests/yr			1995-1997	300 nests/yr	No		Schulz 1987, Groombridge and Luxmoore 1989	Suganuma <i>et al.</i> 1999.
8-c	(Bangka-Belitung Province) Momperang & Pesemut Islands (islands in the vicinity of Belitung)	NN	1980s	3,250 nests/yr	1996	400 nests/yr	2000-2005	350 nests/yr	Yes A, B	270 nests/yr	Schulz 1987, Suganuma <i>et al.</i> 1999.	Suganuma unpublished data, <i>in litt.</i> to J. Mortimer, 2006. <www.elna.or.jp>
	(Bangka-Belitung Province) Kimar Island	NN			1996, 1999	232 nests/yr	2000-2005	~230 ? nests/yr	No <sup>a</sup>			

Index #	Nesting site	Data type	Past estimate 1		Past estimate 2		Recent estimate		Currently protected?		Citation (past)	Citation (recent)
			Years	Mean	Years	Mean	Years	Mean	Yes/No	# Nests		
8-d	(Bangka-Belitung Province) <b>Tiga Islands</b>	NN	1980s	350 nests/yr			1995-1997	150 nests/yr	No		Groombridge and Luxmoore 1989.	Suganuma <i>et al.</i> 1999
8-e	(DKI Jakarta Province) <b>Seribu Islands National Park</b>	NN	1980s	500 nests/yr			since 1995	150 nests/yr	Yes <sup>b</sup> E,D,F	50 nests/yr	Groombridge and Luxmoore 1989.	Suganuma 2005, H. Suganuma <i>in litt.</i> to J. Mortimer (6 Oct 2006).
	Irian Jaya Barat Province <b>Jamursba-Medi region</b>	NN					1999-2005	21 nests/yr	Yes A,B,C,D	21 nests/yr		H. Suganuma <i>in litt.</i> to J. Mortimer (6 Oct 2006).
8-f	(Jawa Tengah Province) <b>Karimunjawa</b>	NN	1980s	300 nests/yr			since 1995	100 nests/yr	No		Groombridge and Luxmoore 1989, Salm 1984.	Suganuma <i>et al.</i> 1999, Suganuma 2005.
8-g	(Jawa Timur Province) <b>Alas Purwo National Park</b>	NN	1983-1989	7.6 nests/yr	1990-1996	8.6 nests/yr	1997-2002	8.7 nests/yr	Yes G	9 nests/yr	Alas Purwo National Park, unpub. data; K. Putra, pers. comm. to J. Mortimer 2006.	Alas Purwo National Park, unpub. data; K. Putra, pers. comm. to J. Mortimer (2006)
8-h	(Jawa Timur Province) <b>Meru Betiri National Park</b>	NN	1980-1989	14.8 nests/yr	1990-1994	3.2 nests/yr	1995	<3 nests/yr	Yes ???	<3 nests/yr	Wetlands International 1997.	Wetlands International 1997.
8-i	(Kalimantan Barat Province) <b>Paloh</b> (4 beaches)	NN	1980s	250 nests/yr			1990-1995	450-478 nests/yr	No		Schulz 1987.	H. Suganuma <i>in litt.</i> to J. Mortimer (2006); Suganuma 2005.
8-j	<b>Kalimantan Selatan Province</b>	NN	1980s	1,000 nests/yr			since 1995	400 nests/yr	No		Groombridge and Luxmoore 1989.	Suganuma 2005.

Index #	Nesting site	Data type	Past estimate 1		Past estimate 2		Recent estimate		Currently protected?		Citation (past)	Citation (recent)
			Years	Mean	Years	Mean	Years	Mean	Yes/No	# Nests		
	(Lampung Province) <b>Segama Besar &amp; Segama Kecil</b>	NN			1996-2000	191 nests/yr	2001-2005	245 nests/yr	Yes	245 nests/yr A, B	Suganuma unpub. data, <i>in litt.</i> to J. Mortimer, 6 Oct 2006. <a href="http://www.elna.or.jp">www.elna.or.jp</a>	Suganuma unpub. data, <i>in litt.</i> to J. Mortimer, 6 Oct 2006; < <a href="http://www.elna.or.jp">www.elna.or.jp</a> >
8-k	(Riau Province) <b>Anambas Islands</b>	NN	1980s	800 nests/yr			2002	300 nests/yr	No		Schulz 1987	Akil <i>et al.</i> 2004.
8-l	(Riau Province) <b>Natuna Besar Islands</b>	NN	1980s	200 nests/yr			2002	50 nests/yr	No		Schulz 1987.	Akil <i>et al.</i> 2004.
8-m	(Riau Province) <b>Tambelan Islands</b>	NN	1980s	1,000 nests/yr	1995-1997	<500 nests/yr	2003	300 nests/yr	No <sup>c</sup>		Schulz 1987.	Suganuma <i>et al.</i> 1999, Akil <i>et al.</i> 2004.
	(Riau Province) <b>reported from 15 beaches</b>	NN	early 1970s	100 nests/beach/yr = ~1,500?	1984-1992	<10 nests/beach/yr = ~150			na		Schulz 1995, <i>in litt.</i> to K. Bjørndal	
8-n	(Sulawesi Selatan Province) <b>Spermonde</b> (Panambungan)	NN	1980S	40 nests/yr			1995-1997	4 nests/yr	No		Schulz 1984.	Suganuma <i>et al.</i> 1999.

Nesting site	Data type	Past estimate 1		Past estimate 2		Recent estimate		Currently protected?	
		Years	Mean	Years	Mean	Years	Mean	Yes/No	# Nests
<b>Population Trends</b> for <u>14 sites</u> for which "Past Estimate 1" and "Recent Estimate" were collected during comparable time periods (~1985 and ~2005)		NN	1980s = (1985)	8,113 nests/yr		1995-2005 = (2005)	2,630 nests/yr		
<b>Current Protected Status</b> of nests at <u>17 sites</u> for which "Recent Estimates" exist									
Protected**								Yes	598
Not Protected**								No	2,528
Total Nests (for <u>17 sites</u> )									3,126

\*\* Only ~19 % of all egg clutches laid at the 17 sites are currently protected.

a Protected by ELNA & ISTRC in 1996, but no protection during 1999-2005 due to occupation by pirates. Survey conducted 2006 by ELNA & ISTRC with plans to protect in 2007.

b Protected by PHPA and Japan Bekko Association in 1995-2000, project continued by National Park Rangers of Seribu Islands.

c On 17 July 2006 legislation passed to eliminate the Forestry Minister Resolution which overrode national protective legislation to allow the collection of turtle eggs in Riau Province (Press Release by Profauna, 22 September 2006). Enforcement will be difficult, however, because so much of the local economy depends on egg collection (H. Sukanuma *in litt.* to J. Mortimer, 10 Mar 2007).

**PAC-Table 5.** Current population estimates and qualitative information about status and trends for reviewed hawksbill populations in the Pacific Ocean. Population estimates are based on nesting females / yr, but where estimates are derived from numbers of nests, a bracketed figure of 3-5 nests per female is used to convert from numbers of nests to numbers of females.

Index #	Locality	Current Population Estimate	Comments (Source)	Status / Trends (Source)
<b>PACIFIC OCEAN: WESTERN</b>				
7	<b>Australia</b> Torres Strait-Northern Great Barrier Reef (GBR) sub-population  Index site = Milman Island	Torres Strait-Northern Great Barrier Reef (GBR) sub-population: = ~4,000 females/yr  <i>Milman Island (Index Site) = ~300-400 females/yr</i>	Within Torres Strait and western Cape York Peninsula, half of all nesting is outside protected habitat (Limpus 2004). On inner shelf of northern GBR, most rookeries are within National Parks; but these nesters are killed on foraging grounds in adjacent countries, particularly Solomon Islands (Limpus 1997, 2004). From ~1850 into the 1930s a ton of tortoiseshell was exported annually from Torres Strait (i.e., an annual take of >1,000 adult Hawksbills) (Limpus, 2004). Since 1968, Hawksbills have been protected in Queensland.	Declining.  Milman Island index population, surveyed since 1990, declining at rate of 3% annually (Limpus 1997, Limpus <i>et al.</i> 1997, Limpus and Miller 2000, K. Dobbs, <i>in litt.</i> to J. Mortimer 2001). If trends continue, projected rate of decline for the Torres Strait-Northern GBR sub-population would be >90% by the year 2020, i.e., in less than one Hawksbill generation (Limpus 2004).
	<b>Australia</b> Northeastern Arnhem Land sub-population	~2,500 females/yr	Most Hawksbill rookeries of Arnhem Land are outside National Parks or other habitat managed for conservation purposes (Limpus 2004). Populations are not regularly surveyed. Shell exports from Australia to Japan since 1950: 29,109 turtles (25,616 kg)	Trends unknown.  Entanglement of juvenile Hawksbills in marine debris including fishing gear is significant threat in northern Australian waters (Kiessling 2003, White 2004).

Index #	Locality	Current Population Estimate	Comments (Source)	Status / Trends (Source)
8	<b>Indonesia</b> <u>Entire</u> Country	6,808-9,077 nests/yr (See also PAC- Table 3)  1,362-3,026 females/yr	<p>Estimate based on assumption that 68 % decline in nesting numbers recorded from the mid-1980s to the present at the 14 sites described in PAC-Table 3, typifies national trend. Based on Schulz's 1987 estimates of 21,000 to 28,000 nests per year during the mid-1980s, a 68% decline translates to 6,808 - 9,077 egg clutches produced annually. Parsons (1972) described "shoal waters of the East Indian archipelago [to] have been the most productive of all the world's seas in tortoise shell". Dammerman (1929) reported much hunting to produce shell equivalent of 160,700 Hawksbills exported to Japan, Singapore, and Netherlands during 1918-1927; but he considered intensive egg collection as greatest danger to species' survival.</p> <p>Shell exports to Japan since 1950: 155,655 turtles (116,741 kg).</p> <p>Shell exports from Singapore to Japan since 1970 probably from Indonesia: 59,215 turtles (44,411 kg).</p> <p>Exports of stuffed juveniles to Japan 1970-1986: 428,859 and 88,539 specimens from Singapore of probable Indonesian origin (Milliken and Tokunaga 1987).</p>	<p>Depleted and declining.</p> <p>Joop Schulz (<i>in litt.</i> to K. Bjorndal, 1995, cited in Meylan and Donnelly 1999) reported: "Almost every egg is taken in virtually every nesting place in Indonesia, however small or far-off it may be" and fishermen complained that Hawksbill had become rare with large sizes seldom caught.</p> <p>Intensive egg collection began at previously uninhabited islands of the Java Sea in 1960-70 when the Bugis people moved from South Sulawesi to avoid civil war (Suganuma <i>et al.</i> 1999). More than 80% of egg clutches are still collected today at surveyed beaches (see PAC-Table 4).</p>

Index #	Locality	Current Population Estimate	Comments (Source)	Status / Trends (Source)
	<b>Japan</b>	rare	Found off southern main islands of Japan; nesting occurs in Ryukyu Archipelago and Ogasawara Islands; considered in danger of extinction since 1985 (Groombridge and Luxmoore 1989). Still fished on foraging grounds (TRAFFIC East Asia-Japan 2000). Shell from Ryukyus since 1950:2171 turtles (1,628 kg). Stuffed juveniles from Ryukyus 1970-1986: 13,438.	Remnant population.
9	<b>Malaysia (East)</b> Sabah Turtle Island	~ 395 nests/yr 80-130 females/yr	Over-exploitation of eggs prior to 1965 when protection began at Turtle Islands Park (de Silva 1982). Likelihood that incubation of all eggs in hatchery since 1965 has feminized offspring (Mortimer 1991c). During 2006, severe erosion of nesting beach habitat at Gulisaan Island was a threat to nesting population (P. Basinthal, <i>in litt.</i> to J. Mortimer, 1 Sep 2006). Shell exported to Japan from Sabah since 1950: 8,089 turtles (6,067 kg).	Probably stable since 1985.
10	<b>Malaysia (West)</b> Terengganu	18 nests/yr 4 - 6 females/yr	Eggs taken in Terengganu since early 20 <sup>th</sup> Century (Siow and Moll 1982), and over-exploitation caused significant decline (Liew 2002). Exports of shell to Japan since 1950 from West Malaysia: 21,344 turtles (16,008 kg) (Records do not specify from which State(s) in West Malaysia the shell originated).	Depleted and declining.

Index #	Locality	Current Population Estimate	Comments (Source)	Status / Trends (Source)
	<b>Papua New Guinea</b>	Low density nesting throughout country? 500-1,000(?) females/yr	<p>Rough estimate provided by B. Krueger (in litt. to J. Mortimer, 27 May 2007) based on 2006-07 surveys:</p> <p><b>a)</b> at Kudube, Takala and Udube Islands, Huon coast, Morobe Province indicate 150-200 nests; but human egg collection and monitor lizard predation allow no more than 10% of nests to hatch;</p> <p><b>b)</b> at Siar Bay, Morobe Province, indicate no nesting at what was previously a sizeable nesting rookery.</p> <p>Heavy exploitation of foraging turtles (&gt;300 annually) reported from Umboi Island, Morobe Province, Manus Island, Manus Province, also subject to large scale take of foraging animals (B. Krueger in litt. to J. Mortimer, 27 May 2007).</p> <p>Spring (1982) reported nesting in East Sepik Province on mainland and islands, in West Sepik Province on islands, on Long island and mainland beaches of Madang Province, on islands of Central Province. Recent surveys conducted at Long Island in Madang Province (Wilson <i>et al.</i> 2004).</p> <p>Shell exports to Japan since 1950: 1,121 turtles (841 kg).</p>	Probably declining



Index #	Locality	Current Population Estimate	Comments (Source)	Status / Trends (Source)
	<b>Philippines</b>	Low density nesting throughout country < 500 females/yr ?	Mindanao coast and Sulu district of southern Philippines historic source of shell (Parsons 1972). According to Seale (1917), outlying islands of the Sulu Archipelago famous for their Hawksbill; in 1917, almost all of 8,000 kg of shell collected each year were exported to Japan. Hawksbill abounded in most areas during the 20 <sup>th</sup> Century but were reduced by 1980 due to heavy exploitation (de Celis 1982). Shell exports to Japan since 1950: 51,259 turtles (38,444 kg). Exports of stuffed juveniles to Japan 1970-1986: 8,698.	Depleted and declining. Population decline due to exploitation for shell, meat and eggs (Alcala 1980 as cited in Groombridge and Luxmoore 1989, de Celis 1982, Groombridge and Luxmoore 1989, Palma 1997). In 1980, Alcala reported that virtually every nesting turtle was killed in the Central Visayas, and believed the same occurred throughout Philippines (Meylan and Donnelly 1999).
12	<b>Thailand</b> Gulf of Thailand	~ 20 females/yr	Ko Khram is Thailand's most important <i>Hawksbill</i> nesting site, controlled and protected by Royal Thai Navy since 1950s. During 1980s, most of eggs laid at Ko Khram and nearby islands sold to Navy Officers and remainder incubated in hatchery, and all hatchlings head-started at Ko Man Nai (Mortimer 1988). Other threats include: mortality from heavy trawl activity and poaching of nesting females (Polunin 1977 as cited in Mortimer 1988, Mortimer 1988); disturbance from bright lights and noise from jetty built in 1970s (Polunin 1977 as cited in Mortimer 1988). Shell exports to Japan since 1950: 27 turtles (20 kg).	Depleted and declining. Serious decline in nesting activity since 1950s, including: recorded decline of 43% during 1973-2005; and estimated decline of 75% during 1956-2005 (Polunin and Nuija 1982, Charuchinda and Monanunsap 1998, M. Charachinda, unpubl. data). Nesting at other sites in Gulf of Thailand now insignificant (Charuchinda and Monanunsap 1998).

Index #	Locality	Current Population Estimate	Comments (Source)	Status / Trends (Source)
	<b>Vietnam</b>	~100 females/yr	Despite declines in the nesting population, Vietnam still has “a strong continuing local tortoiseshell industry” (Le Dien and Broad, 1995; N. Pilcher, <i>in litt</i> to J. Mortimer, 2002) and international export trade. Hawksbills are also slaughtered for meat (Hamann <i>et al.</i> , 2006). In 2002, Vietnam instituted full domestic and trade protection status to Hawksbills (Hamann <i>et al.</i> 2006).	Depleted and probably declining Sixty-five years ago Hawksbill were common along the coast of Vietnam (Bourret 1941 cited in Groombridge and Luxmoore 1989). Egg collection at Cochin China reduced Hawksbill populations by 1923 (Le Poulain 1941 cited in Groombridge and Luxmoore 1989)
<b>PACIFIC OCEAN: CENTRAL</b>				
	<b>American Samoa and Western Samoa</b>	<10-30 females/yr	Estimate based on Tuato'o-Bartley <i>et al.</i> (1993), Grant <i>et al.</i> (1997), G. Balazs, <i>in litt.</i> to J. Mortimer, 15 May 2007)	Depleted and declining.

Index #	Locality	Current Population Estimate	Comments (Source)	Status / Trends (Source)
	<b><i>Fiji</i></b>	~500 plus nests / yr  100-200 females/yr	Estimate based on Batibasaga (2002) and K.T. MacKay (in litt. to J. Mortimer, 15 May 2007).  Over-exploited for more than 100 years (Troëng, 1996). In latter half of the 20 <sup>th</sup> Century intense local exploitation of eggs and adults for food and a major shell carving industry (Groombridge and Luxmoore 1989). Shell exports banned in 1991 (Daly 1991). Increased awareness in young people (Troëng 1996).  Domestic tourist trade in tortoiseshell curios and whole carapaces continues with suggestions of an underground export trade via Asian fishing boats (Laveti and MacKay, MS in prep., cited in K.T. MacKay in litt. to J. Mortimer, 15 May 2007).  Shell exports to Japan 1950-1990: 14,489 turtles (12,751 kg).	Depleted.  Decline of 50% in 20 years reported at Namenalala, a major <i>Hawksbill</i> rookery hosting 30-40 females/yr (K. MacKay pers. comm. to J. Mortimer 23 Feb 2007).  On Tavarua Island, in 1944, Mr. Sadolo reported >100 Hawksbills on a single night in March-April (Sovaki, 1997, cited by K. MacKay in litt. to J. Mortimer, 15 May 2007); but during two nights on Tavarua Island (13 Nov and 4 Dec, 1970) 6 recent nests, but no turtles were reported (Hirth 1971, cited by K. MacKay in litt. to J. Mortimer, 15 May 2007).
	<b><i>Guam</i></b>	5-10 females/yr	Based on rough estimate from G. Davis (NMFS), in litt. to J. Mortimer, 15 May 2007	Depleted and declining.
	<b><i>Hawaii</i></b>	5-10 females/yr	Nest counts and in-water observations indicate this depleted population is increasing (G. Balazs (NMFS), in litt. to J. Mortimer, 15 May 2007)	Depleted and increasing.
	<b><i>Micronesia</i></b> Entire area	~300 females/yr	Nesting is sparsely distributed among thousands of islands and atolls including the Palau Republic discussed below (NMFS and USFWS 1998)	Depleted and declining.

Index #	Locality	Current Population Estimate	Comments (Source)	Status / Trends (Source)
	<b>Palau Republic</b>	20-50 females/yr	Comprises largest nesting population in Micronesia (NMFS and USFWS, 1998). Exploited for meat, eggs, and shell for local consumption (Meylan and Donnelly, 1999).	Depleted and declining.
11	<b>Solomon Islands</b>	800-1,200 nests/yr  200-300 females /yr	Estimated nests/yr based on Wilson <i>et al.</i> (2004), except where otherwise noted: Arnavon Islands and adjacent sites, 500-600 (Ramohia and Pita 1996); Shortlands (100-200), Marovo (50), Ramos (50), Santa Cruz (50-200), and Russell (50-100). Long history of intense shell trade. In mid-1990s, most females killed in first nesting season, with 90% of breeders being first time nesters (Meylan and Donnelly 1999). Protection in the Arnavon Islands Community Marine Conservation Area (ACMCA) significantly improved survival of nesters (J. Pita and P. Ramohia, pers. comm. to J. Mortimer; J. Mortimer, 2002) Shell exports to Japan 1950-1990: 39,090 turtles (34,399 kg).	Depleted and declining. In second half of the 20 <sup>th</sup> Century nesting numbers declined throughout the Solomons (Groombridge and Luxmoore 1989) to ~ 500 females/yr. From 1988 to 1997, Limpus (1997) estimated population declines of more than 50%. Current subsistence take is unsustainable (Broderick and Pita 2005). In communities adjacent to Arnavon Community Marine Conservation Area (ACMCA) estimated 825 Hawksbill of all sizes slaughtered/yr (Broderick and Pita 2005). Of captured Hawksbills tagged and released by Broderick in adjacent communities, 30% were recaptured, and half of those subsequently slaughtered (Broderick and Pita, 2005).

Index #	Locality	Current Population Estimate	Comments (Source)	Status / Trends (Source)
	<b>Vanuatu</b>	>300 females /yr	Scattered nesting throughout the country, especially at: a) Banks/Torres; b) Malekula; c) Epi, Green; and d) Aneityum (Wilson <i>et al.</i> 2004). Surveys in 2006-07 (K. MacKay in litt. to J. Mortimer, 15 May 2007) identified two beaches: a) Moso Island (Efate) with (>100 nests); and b) Bamboo Bay (Malakula) (>200 nests). The nesters are larger than reported elsewhere (mean CCL=94 cm; including some with CCL >100 cm). Feral dog predation a problem at some sites.	Probably declining. Subject to heavy exploitation at some sites (i.e., Malekula) while little or no pressure at others (Wilson <i>et al.</i> 2004). Recently less exploited in many areas (esp. foraging populations) (K. MacKay, pers. comm. to J. Mortimer, 23 Feb 2007) due to public awareness programmes (Petro. 2002).
<b>PACIFIC OCEAN: EASTERN</b>				
	<b>El Salvador</b>	70 females/yr	In 2007, 72 Hawksbills reported on 3 beaches (C. Hasbun pers. comm. to M. Donnelly, 2008).	Remnant population
13	<b>Mexico</b>	Low nesting numbers <15 females/yr	Low nesting numbers in Jalisco, Nayarit, and Tres Marias Islands (Seminoff <i>et al.</i> , 2003b). Once found abundantly along the eastern Pacific coast but now very rare (Cliffon <i>et al.</i> , 1982; Seminoff <i>et al.</i> , 2003b). Documented hybridization between Hawksbill and Green Turtles a cause for concern (Seminoff <i>et al.</i> , 2003a). Occurrence of immatures in Gulf of California indicate nesting nearby; Baja California should be a priority for regional recovery efforts (Seminoff <i>et al.</i> , 2003b).	Remnant populations

**ATL-Table 1.** Atlantic Ocean localities of importance to *Eretmochelys imbricata* (n= 34), including 12 Index Sites (and their assigned reference numbers), for which quantitative data exist on past and present abundance (ATL-Table 2). Long-term changes in population size were calculated with these data and are presented in ATL-Tables 3-6 and ATL-Figure 1. ATL-Table 7 presents current status and qualitative data pertaining to population trends for all 34 sites. Locations of the 12 Index Sites are shown on the map in W-Figure 1.

Index #	Nesting Sites	ATL-Table(s)	Justification
<b>ATLANTIC OCEAN: INSULAR CARIBBEAN</b>			
	Antigua / Barbuda (general)	7	
14	Antigua: Jumby Bay	2, 5, 6, 7	Monitored since 1987 & well-protected
15	Bahamas	2, 3, 5, 6, 7	Historic shell trade stats
16	Barbados	2, 6, 7	Monitored since mid-1980s & well-protected
	British Virgin Islands	7	
17	Cuba (Doce Leguas Cays)	2, 4, 5, 6, 7	Historical records and recent monitoring
	Dominican Republic	7	
	French West Indies: Guadeloupean Archipelago	7	
	French West Indies: Martinique	7	
	Jamaica	7	
	Grenada	7	
	Puerto Rico: Culebra, Caja de Muertos, & Humacao	7	
18	Puerto Rico: Mona Island	2, 5, 6, 7	Long term monitoring & protection since 1970s
	St. Kitts	7	
	Trinidad & Tobago	7	
19	US Virgin Islands: Buck Island Reef Nat'l Monument	2, 5, 6, 7	Long term monitoring & protection
	US Virgin Islands: Sites outside Buck Island National Monument	7	
<b>ATLANTIC OCEAN: WESTERN CARIBBEAN MAINLAND</b>			
	Belize: Manatee Bar, Sapodilla Cays, South Water Cay	7	
	Colombia: Isla Fuerte	7	
	Colombia: San Andres Archipelago	7	

Index #	Nesting Sites	ATL-Table(s)	Justification
20	Costa Rica: Tortuguero National Park	2, 3, 6, 7	Long term monitoring & protection since 1956
	Costa Rica: Cahuita	7	
	Honduras: Bay Islands	7	
21a	Mexico: Yucatan Peninsula	2, 7	Long term monitoring since late 1970s
21b	Mexico: Campeche State	2, 5, 6, 7	Long term monitoring since late 1970s
22	Nicaragua: El Cocal	2, 3, 6, 7	Monitored in the 1970s and early 2000s
	Nicaragua: Miskito Coast	7	
	Nicaragua: Pearl Cays	7	
	Panama: Bastimentos Island National Marine Park	7	
23	Panama: Chiriqui Beach	2, 3, 6, 7	Historical records and recent monitoring
	Venezuela: Los Roques & Paria region	7	
<b>ATLANTIC OCEAN: SOUTH WESTERN</b>			
24	Brazil	2, 3, 6, 7	Long term monitoring since 1980
<b>ATLANTIC OCEAN: EASTERN</b>			
25	Equatorial Guinea (Bioko)	2, 3, 6, 7	Historical records and recent monitoring
	São Tomé and Príncipe	7	

**ATL-Table 2.** Quantitative evaluation of nesting activity and population trends in the Atlantic Ocean based on available *Past* and *Recent* estimates for *Eretmochelys imbricata* at 12 sites. Data codes include: NF, numbers of nesting females; NN, numbers of nests; SNF numbers of slaughtered nesting females; TSE, Tortoiseshell Export Statistics; and UPE, unit patrol effort at the nesting beach. A bracketed figure of 3-5 nests per female was used to convert from number of nests to numbers of females, unless source data reported numbers of females. All values are based on annual means unless otherwise stated.

Index #	Index Nesting Site	Data type	Past Estimate 1		Past Estimate 2		Recent Estimate		Citation (Past)	Citation (Recent)	
			Years	Mean	Years	Mean	Years	Mean			
ATLANTIC OCEAN: INSULAR CARIBBEAN											
14	Antigua Jumby Bay	NF	1987-1991	29 females/yr	1997-2001	33 females/yr	2002-2005	52 females/yr	Richardson <i>et al.</i> 1999.	Parish and Goodman 2006; McIntosh <i>et al.</i> , 2003; Stapleton and Stapleton 2004, 2006	
15	Bahamas	TSE	1891-1900	4,186 kg/yr	1932 - 1938	2,055 kg/yr	1970 - 1979	734 kg/yr*	Northcroft, 1900 (cited in McClenachan <i>et al.</i> 2006); export Statistics listed in Commonwealth of Bahamas, Colonial Reports,1932-1964 (Seminoff and Bjorndal unpublished summary, Pandolfi <i>et al.</i> 2003)	Japanese Customs Statistics (1950-1986) (Milliken and Tokunaga 1987, Groombridge and Luxmoore 1989)	
		NN					2006	500-1,000 nests/yr			K. Bjorndal ( <i>in litt.</i> to J. Mortimer, 5 Nov 2006)
		NF						100-333 females/yr			



Index #	Index Nesting Site	Data type	Past Estimate 1		Past Estimate 2		Recent Estimate		Citation (Past)	Citation (Recent)
			Years	Mean	Years	Mean	Years	Mean		
16	<b>Barbados</b>	NF	mid-1980s	60 females/yr	1997-1998	103 females/yr	2003-2005	483 females/yr	Estimate based on public reports, beach surveys, tagging program (Horrocks 1992, J. Horrocks pers. comm. to A. Meylan, Meylan 1999, Horrocks and Krueger unpub. data	Beggs <i>et al.</i> 2007, Horrocks and Krueger unpub. data
17	Cuba <b>Doce Leguas Cays (all)</b>	NN	1880s	<b>6,000-10,000 nests/ yr</b>			2002	<b>2,000-2,500 nests/yr</b>	<b>Thousands caught annually on nesting beaches (Ballou, 1888, cited in McClenachan 2006)</b>	Cuban Turtle Group <i>in litt.</i> to A. Abreu, Feb 2002
		NF		<b>3,000 females/yr</b>				400-833 females/yr		
	<b>Doce Leguas Cays 9 Index Beaches</b>	NN	1997-1999	<b>45.3 nests/yr</b>	2000-2002	<b>52.7 nests/yr</b>	2003-2005	<b>65 nests/yr</b>	<b>F. Moncada, <i>in litt.</i> to M. Donnelly, 10 Nov 2006</b>	F. Moncada, <i>in litt.</i> to M. Donnelly, 10 Nov 2006
		NF		<b>9-15 females/yr</b>		<b>10.5-17.6 females/yr</b>		<b>13-21.7 females/yr</b>		
18	<b>Puerto Rico Mona Island</b>	NN	1974, 1984-1990	117 nests/yr	1991-1998	297 nests/yr	1999-2005	742 nests/yr	Unpubl. data, C.E. Diez <i>in litt.</i> to J. Mortimer, 2006	Unpubl. data, R.P. van Dam and C.E. Diez (C.E. Diez <i>in litt.</i> to J. Mortimer, 2006)
		NF		23-39 females/yr		59 - 99 females/yr		148 - 247 females/yr		

Index #	Index Nesting Site	Data type	Past Estimate 1		Past Estimate 2		Recent Estimate		Citation (Past)	Citation (Recent)
			Years	Mean	Years	Mean	Years	Mean		
19	US Virgin Islands <b>Buck Island Reef Nat'l Monument</b> (St. Croix)	NF	1988-1994	23 females/yr	1995-2000	26 females/yr	2001-2006	56 females/yr	Z. Hillis-Starr and B. Phillips ( <i>in litt.</i> to J. Mortimer 2006).	Unpubl. data (Z.Hillis- Starr, <i>in litt.</i> to J. Mortimer 2006)
<b>ATLANTIC OCEAN: WESTERN CARIBBEAN MAINLAND</b>										
20	Costa Rica <b>Tortuguero National Park</b>	UPE	1956-1960	0.0606 tracks/night/km	1986-1990	0.0140 tracks/night/km	2001-2005	0.0116 tracks/night/km	Caribbean Conservation Corporation, unpublished data; Troëng <i>et al.</i> 2005	Caribbean Conservation Corporation, unpublished data; Troëng <i>et al.</i> 2005
		NF					2005	~ 10 females/yr		Caribbean Conservation Corporation, unpublished data
21a	Mexico <b>Entire Yucatan Peninsula</b> (including Campeche, Yucatan and Quintana Roo)	NN	1990-1995	1,614 nests/yr	1996-2001	4,148 nests/yr	2001-2006	2,672 nests/yr	Abreu-Grobois <i>et al.</i> , 2005; Abreu-Groboi <i>in litt.</i> to J. Mortimer, 9 Feb 2007. Based on unpublished data collected in: <b>a)</b> Yucatán and Quintana Roo by: Pronatura Península de Yucatán, SEMARNAT, CONANP, Secretaría de Ecología de Yucatán; and <b>b)</b> Campeche by: Conanp-APFFLT, SEMAR V Zona Naval, Secretaria de Ecología Gob. del Estado, Enlaces con tu Entorno AC, Marea Azul AC, Desarrollo Ecologico Cd. del Carmen AC, Quelonios AC, UNACAR, Universidad Autónoma de Campeche, H. Ayuntamiento del Carmen, Pronatura PPY, Profepa.	
		NF		322-538 females/yr		830-1,383 females/yr		534-891 females/yr		

Index #	Index Nesting Site	Data type	Past Estimate 1		Past Estimate 2		Recent Estimate		Citation (Past)	Citation (Recent)
			Years	Mean	Years	Mean	Years	Mean		
21b	Mexico <b>Campeche State</b> (Yucatan Peninsula)	NN	1985-1986, 1990-1991	388 nests/yr	1992-1998	1,748 nests/yr	1999-2005	2,236 nests/yr	Márquez <i>et al.</i> , 1987; Vicente Guzmán pers. comm. to A. Abreu-Grobois 2006, Garduño <i>et al.</i> 1999, M. Medina pers. comm. to A. Abreu-Grobois 2006	Vicente Guzmán, pers. comm. to A. Abreu-Grobois 2006, Garduño <i>et al.</i> 1999, M. Medina pers. comm. to A. Abreu-Grobois 2006
		NF		78-129 females/yr		350-583 females/yr		447-745 females/yr		
22	Nicaragua <b>El Cocal</b> (near San Juan del Norte)	NN	1970s	~ > 300 nests /yr			2000	75 nests/yr	Lagueux and Campbell 2005	Lagueux and Campbell 2005
		NF		60-100 females/yr				15 -25 females/yr		
23	Panama <b>Chiriqui Beach:</b> one mile of total 15 miles	SNF / UPE	early 1950s	35-50 females/mile/ night (peak season)	1980s	1-5 females/mile/ night (peak season)	2003-2005	1-3 females/mile/ night (peak season)	Carr 1956, Carr <i>et al.</i> 1982, Meylan and Donnelly 1999	Ordoñez, pers. comm. to A. Meylan
		NN					2003-2005	421 nests/yr		Meylan <i>et al.</i> 2006
		NF						84-140 females/yr		

Index #	Index Nesting Site	Data type	Past Estimate 1		Past Estimate 2		Recent Estimate		Citation (Past)	Citation (Recent)
			Years	Mean	Years	Mean	Years	Mean		
ATLANTIC OCEAN: SOUTH WESTERN										
24	Brazil	NN	1901	>8,750 nests/yr			2005	~1,750 nests/yr	N. Marcovaldi, pers. comm. to J. Mortimer 2006	Marcovaldi <i>et al.</i> in press
		NF		1,750-2,917 females/yr				350-585 females/yr		
ATLANTIC OCEAN: EASTERN										
25	Equatorial Guinea Bioko	UPE	1940s	200-300 females (all species) per night at peak season	1980s	50-100 females (all species) per night at peak season	1996-2005	7 females/yr	T. Butynski in litt. to K. Bjorndal, 20 April 1986, cited in Groombridge and Luxmoore, 1989	Tomás <i>et al.</i> 2000, Rader <i>et al.</i> 2006

**ATL-Table 3.** Summary of estimated population change over 3 generations for 6 Atlantic Ocean Index Sites for which there are historical data prior to the 1980s. Figures derive from the Past and Recent Estimates presented in ATL-Table 2, and from Exponential and Linear extrapolation functions (IUCN 2001a). Extrapolation functions are used only when there is a suspected change in the subpopulation size over a specific time interval outside of the period represented by data in ATL-Table 2. Where bracketed estimates are presented in ATL-Table 2, the mid-point is used here. In such cases, unless otherwise noted, both linear (L) and exponential (E) functions are used due to a lack of information on the true rate of change over the time interval. All values are based on annual means.

Index #	Index Sites <u>WITH</u> Historical Data Prior to the 1980s	Raw Data (from ATL-Table 2)		Notes on Population Trajectories & Comments on Current Status	Past Annual Nesting Female Subpopulation Size (3 generations back)	Present Annual Nesting Female Subpopulation Size (2005)	% Change over 3 generations
		Past	Present				
ATLANTIC OCEAN: INSULAR CARIBBEAN							
15	Bahamas	Proxy:		1901-1979: Declining (McClenachan <i>et al.</i> 2006; Export Statistics for Commonwealth of Bahamas, Colonial Reports, 1932-1964 (Seminoff and Bjorndal, unpubl. summ.); Japanese Customs Statistics, 1950-1986, compiled by Groombridge and Luxmoore 1989.	Proxy:		
		4,186 kg shell exported/year (1891-1900)	734 kg shell exported/ year (1970- 1979)		3,734 kg (L) (1901)	164 kg (L) (1979)	- 96%
					3,458 kg (E) (1901)	380 kg (E) (1979)	-89%
		Population estimate:		1979-2005: Continued populations decline likely due to poaching for meat & habitat destruction (K. Bjorndal, <i>in litt.</i> to J. Mortimer, 5 Nov 2006) Declining (B.Riegl, <i>in litt.</i> to J. Mortimer, 9 Oct 2006)	Extrapolated population estimate:		
		na	217 females/yr (2005)		5,425 (L) females/yr (1901)	217 females/yr (2005)	- 96%
				1,972 (E) females/yr (1901)	217 females/yr (2005)	-89%	

Index #	Index Sites <u>WITH</u> Historical Data Prior to the 1980s	Raw Data (from ATL-Table 2)		Notes on Population Trajectories & Comments on Current Status	Past Annual Nesting Female Subpopulation Size (3 generations back)	Present Annual Nesting Female Subpopulation Size (2005)	% Change over 3 generations
		Past	Present				
ATLANTIC OCEAN: WESTERN CARIBBEAN MAINLAND							
20	Costa Rica Tortuguero National Park	Proxy:  0.0606 tracks/night/km (1956-1960)	0.0116 tracks/night/km (2001-2005)	1901-1955: Trend unknown (assumed stable)  1956-2005: Declining (Carr and Stancyk 1975, Bjorndal <i>et al.</i> 1993, Troëng <i>et al.</i> 2005, Caribbean Conservation Corporation unpubl. data)	Proxy:  0.0606 tracks/night/ km (1956)	0.0116 tracks/night/km	-81%
		Population estimate:  na	10 females/yr (2005)		Population estimate:  52 females/yr (1956)	10 females/yr (2005)	-81%
22		Nicaragua El Cocal (near San Juan del Norte)	80 females/yr (1970s)	20 females/yr (2000)	1901-1970s: Declining (Nietschmann 1973)  1970s-2005: Declining (Lagueux and Campbell 2005)	258 females/yr (L) (1901)	8 females/yr (L) (2005)
	4,536 females/yr (E) (1901)					14 females/yr (E) (2005)	-99%

Index #	Index Sites <u>WITH</u> Historical Data Prior to the 1980s	Raw Data (from ATL-Table 2)		Notes on Population Trajectories & Comments on Current Status	Past Annual Nesting Female Subpopulation Size (3 generations back)	Present Annual Nesting Female Subpopulation Size (2005)	% Change over 3 generations
		Past	Present				
23	Panama <b>Chiriqui Beach:</b> <i>one mile out of 15 miles</i>	Proxy:  <b>42.5</b> females/mile/ night (peak season) (early 1950s)  Population estimate:  <b>na</b>	<b>2</b> females/mile/ night (peak season) (2003-2005)  <b>112</b> females / yr (2005)	<b>1901-1950s:</b> Trend unknown <b>1950s-1980s:</b> In steep decline (Carr 1956, Carr <i>et al.</i> 1982, Meylan and Donnelly 1999, Ordoñez pers. comm. to A. Meylan) <b>1980s-2005:</b> Declining (Meylan <i>et al.</i> 2006)	Proxy:  <b>42.5</b> females/ mile/ night (peak season) (early 1950s)  Extrapolated population estimate:  <b>2,380</b> females / yr (early 1950s)	<b>2</b> females/ mile/ night (peak season) (2003-2005)  <b>112</b> females / yr (2005)	<b>-95 %</b>     <b>-95 %</b>
ATLANTIC OCEAN: SOUTH WESTERN							
24	<b>Brazil</b>	<b>2,333</b> females/yr (1901)	<b>468</b> females/yr (2005)	<b>1901-1982:</b> Declined by >-80% due to habitat destruction, directed take, fisheries related mortality (N. Marcovaldi, pers. comm. to J. Mortimer, 2006; 2005; Marcovaldi <i>et al.</i> in press). <b>1982-2005:</b> Increasing (Marcovaldi <i>et al.</i> in press).	<b>2,338</b> females/yr (1901)	<b>468</b> females/yr (2005)	<b>-80%</b>

Index #	Index Sites <u>WITH</u> Historical Data Prior to the 1980s	Raw Data (from ATL-Table 2)		Notes on Population Trajectories & Comments on Current Status	Past Annual Nesting Female Subpopulation Size (3 generations back)	Present Annual Nesting Female Subpopulation Size (2005)	% Change over 3 generations
		Past	Present				
ATLANTIC OCEAN: EASTERN							
25	Equatorial Guinea <i>Bioko</i>	333 females/yr (1945)  100 females/yr (1985)  Assume hawksbills comprise 13% of all turtles, and peak season is 10 days.	7 females/yr (1998)	1901-1940: Trend unknown 1940-present: Intense exploitation of nesting turtles (all species) including hawksbills for meat, shell and eggs. At peak season, 200-300 (in 1940s), and 50-100 (in mid-1980s) turtles taken per night (all species), with "a significant portion" of these being hawksbills (Groombridge and Luxmoore 1989). Average of 7 hawksbills/ yr in 1997-98 (Tomás <i>et al.</i> 2000). Only 4-7 nests recorded each year during 2001-05 (Rader <i>et al.</i> 2006).	360 females/yr (L) (1940)   752 females/yr (E) (1940)	0 females/yr (L) (2005)   4 females/yr (E) (2005)	100 %       -99 %



**ATL-Table 4.** Summary of estimated population change over 3 generations for the Doce Leguas Cays of Cuba. Figures for 1901 and 1985 derive from the Past and Recent Estimates presented in ATL-Table 2, and from Exponential and Linear extrapolation functions (IUCN 2001a). The derivation of figures for 2005 is shown in ATL-Table 5. Extrapolation functions are used where there is a suspected change in the subpopulation size over a specific time interval outside of the period represented by data in ATL-Table 2. Where bracketed estimates are presented in ATL-Table 2, the mid-point is used here. In such cases, unless otherwise noted, both linear (L) and exponential (E) functions are used due to a lack of information on the true rate of change over the time interval. All values are based on annual means.

Index #	Index Sites	Raw Data (from ATL-Table 2)		Notes on Population Trajectories & Comments on Current Status	Estimated Past Annual Nesting Female Subpopulation Size (3 generations back)	Estimated Annual Nesting Female Subpopulation Size (1985)	Estimated Present Annual Nesting Female Subpopulation Size (2005)
		Past	Present				
ATLANTIC OCEAN: INSULAR CARIBBEAN							
17	Cuba Doce Leguas Cays	3,000 females/yr (1880s) 'thousands of females killed annually on the beaches'	617 females/yr (2002)	1901-1992: Declining (McClenachan <i>et al.</i> 2006). 1993-2002: Annual legal foraging ground take of 5,000 reduced to 3,000 in 1993, 1,000 in 1994, and 500 from 1995 onwards (Carrillo <i>et al.</i> 1999). Evidence of increase from 1997-2005 (see ATL-Table 5).	2,673 females/yr (L) (1901)  2,146 females/yr (E) (1901)	963 females/yr (L) (1985)  690 females/yr (E) (1985)	1,178 females/yr (L) (2005)  963 females/yr (E) (2005)

**ATL-Table 5.** Estimated population change for 6 Atlantic Ocean Index Sites that have recorded increases in nesting populations since 1985, but for which no historical data exist prior to the 1980s. Figures derive from the Past and Recent Estimates presented in ATL-Table 2; where these are bracketed estimates only the mid-point is used here. All values are based on annual means. Extrapolated population trajectories between 1901 and 1985 are presented in ATL-Table 6.

Index #	Index Sites	Raw Data (from ATL-Table 2)		Notes on Population Trajectories & Comments on Current Status	Change in Population Size Since Protection Implemented
		Past	Present		
ATLANTIC OCEAN: INSULAR CARIBBEAN					
14	Antigua Jumby Bay	29 females/yr (1987-1991)	52 females/yr (2002-2005)	1901-1987: Declining (Fuller <i>et al.</i> 1992, Meylan 1999) 1987-2001: Stable 2002-2005: Increasing (Depleted but <u>Increasing</u> )	+ 79 % during 19 years  + 23 females
16	Barbados	60 females/yr  (mid-1980s)	483 females/yr  (2003-2005)	1901-1985: Declining (Horrocks 1992, J. Horrocks pers. comm. to A. Meylan, Meylan 1999) 1985-1997: Unknown 1997-2005: Nesting activity increasing (Beggs <i>et al.</i> 2007, Horrocks and Krueger unpublished data. (Depleted but <u>Increasing</u> )	+ 705 % during 20 years  + 423 females
17	Cuba  Doce Leguas Cays	Proxy (9 index beaches at Doce Leguas):  11 females/yr (1997)  Extrapolated Population Estimates:  718 (L) females/yr (1997)  587 (E) females/yr (1997)	18 females/yr (2005)  assuming 64% increase  1,178 (L) females/yr (2005)  assuming 64% increase  963 (E) females/yr (2005)  assuming 64% increase	1901-1992: Declining (McClenachan <i>et al.</i> 2006). 1993-2005: Annual legal take on foraging grounds of 5,000 reduced to 3,000 in 1993, 1,000 in 1994, and 500 from 1995 onwards (Carrillo <i>et al.</i> 1999). Reportedly increasing.  (Depleted but <u>Increasing</u> )	Proxy (9 index beaches at Doce Leguas):  + 64 % during 8 years  Extrapolated Population Estimates:  + 460 (L) females/yr  + 376 (E) females/yr  +418 females

Index #	Index Sites	Raw Data (from ATL-Table 2)		Notes on Population Trajectories & Comments on Current Status	Change in Population Size Since Protection Implemented
		Past	Present		
18	Puerto Rico <i>Mona Island</i>	<b>31</b> females/yr (1974, 1984- 1990)	<b>198</b> females/yr (1999-2005)	<b>1901-1960s:</b> Trend unknown <b>1960s-early 1990s:</b> Declining <b>Early 1990s-2005:</b> Increasing.  (Depleted but <u>Increasing</u> )	<b>+ 539 %</b> during <b>31</b> years  <b>+167 females</b>
19	US Virgin Islands <i>Buck Island Reef Nat'l Monument</i> (St. Croix)	<b>23</b> females/yr (1988-1994)	<b>56</b> females/yr (2001-2006)	<b>1901-1960s:</b> Trend unknown <b>1960s-early 1987:</b> Declining <b>1988-2000:</b> Apparently stable <b>2001-2006:</b> Increasing  (Depleted but <u>Increasing</u> )	<b>+ 143 %</b> during <b>18</b> years  <b>+ 33 females</b>
<b>ATLANTIC OCEAN: WESTERN CARIBBEAN MAINLAND</b>					
21	Mexico <i>Campeche State (Yucatán Peninsula)</i>	<b>104</b> females/yr (1985-1991)	<b>596</b> females/yr (1998-2005)	<b>1901-1977:</b> Declining <b>1978:</b> Protection begun. <b>1985-1999:</b> Increased dramatically (~475%) probably due to local and regional protection (Garduño- Andrade <i>et al.</i> 1999). <b>1999-2004:</b> Declined by -63% in 5 years (Abreu-Grobois <i>et al.</i> 2005) <b>2004:</b> Stopped declining; lowest records for the region. <b>2005-2006:</b> Starting new increase	<b>+473 %</b> during <b>21</b> years  <b>+492 females</b>
<b>Total change: average females/yr</b>				<b>Estimated change in average number nesting per year since 1985</b>	<b>+1,252 females</b>

**ATL-Table 6.** Summary of estimated population change over 3 generations for 12 Atlantic Index Sites including the following: a) Caribbean sites with historic data prior to the 1980s; b) Caribbean sites lacking historic data prior to the 1980s; c) South Western & Eastern Atlantic sites; and d) Regional summaries of all index sites from: i. the Entire Caribbean; and ii. the Entire Atlantic Ocean. Data codes include: R = raw data or figures calculated arithmetically; L= figures calculated from linear extrapolation; and E=figures calculated from exponential extrapolated.

ATL-Table 6 (a). Caribbean sites <u>with</u> historic data prior to the 1980s													
Index #	Index Sites <u>WITH</u> Historical Data Prior to the 1980s	Population Estimate: mean # females/ yr			Estimated Change in Population								
		1901		1985		2005		1901-1985		1985-2005		1901-2005	
		1901		1985		2005	%	# Fem	%	# Fem	%	# Fem	
ATLANTIC OCEAN: CARIBBEAN SITES													
15	Bahamas	L	5,425	L	1,218	R	217	-77.5 %	-4,207	-82.2 %	-1,001	-96.0 %	-5,208
		E	1,972	E	371	R	217	-81.2 %	-1,601	-41.5 %	-154	-89.0 %	-1,755
20	Costa Rica (Tortuguero National Park)	R	52	R	14.3	R	10	-72.5 %	-38	-30.1 %	-4	-80.8 %	-42
17	Cuba (Doce Leguas Cays)	L	2,673	L	963	L	1,178	-64.0 %	-1,710	22.3 %	+215	-55.9 %	-1,495
		E	2,146	E	690	E	963	-67.8 %	-1,456	39.6 %	+273	-55.1 %	-1,183
22	Nicaragua (El Cocal)	L	258	L	56	L	8	-78.3 %	-202	-85.7 %	-48	-96.9 %	-250
		E	4,536	E	43	E	14	-99.1 %	-4,493	-67.4 %	-29	-99.7 %	-4,522
23	Panama (Chiriqui Beach)	L	2,380	L	1,019	R	112	-57.2 %	-1,361	-89.0 %	-907	95.3 %	-2,268
		E	2,657	E.	425			-84.0 %	-2,232	-73.6 %	-313	95.8 %	-2,545
CARIBBEAN INDEX SITES <u>WITH</u> <u>HISTORICAL</u> DATA													
	<u>LINEAR</u> & RAW DATA	L	10,788	L	3,270	L	1,525	-69.7 %	-7,518	-53.4 %	-1,745	-85.9 %	-9,263
	<u>EXPONENTIAL</u> & RAW DATA	E	11,363	E	1,543	E	1316	-86.4 %	-9,820	-14.7 %	-227	-88.4 %	-10,047

ATL-Table 6 (b). Caribbean sites lacking historic data prior to the 1980s

Index #	Index Sites <u>LACKING</u> Historical Data Prior to the 1980s	Population Estimate: mean # females/ yr Note: 1901 figures extrapolated from average population trajectory for 1901-1985 calculated above in ATL-Table 6a.			Estimated Change in Population										
					1901-1985		1985-2005		1901-2005						
		1901	1985	2005	%	# Fem	%	# Fem	%	# Fem					
ATLANTIC OCEAN: CARIBBEAN SITES															
14	Antigua (Jumby Bay)	L	96	R	29	R	52	-69.7 %	-67	+79 %	+23	-45.8 %	-44		
		E	213					-86.4 %	-184			-75.6 %	-161		
16	Barbados	L	198	R	60	R	483	-69.7 %	-138	+705 %	+423	+143.9%	+285		
		E	441					-86.4 %	-381			+9.5 %	+42		
21b	Mexico (Campeche)	L	343	R	104	R	596	-69.7 %	-239	+473 %	+492	+73.8 %	+253		
		E	765					-86.4 %	-661			- 22.1 %	-169		
18	Puerto Rico (Mona)	L	102	R	31	R	198	-69.7 %	-71	+539 %	+167	+94.1 %	+96		
		E	228					-86.4 %	-197			-13.2 %	-30		
19	U.S. Virgin Islands (Buck Island)	L	76	R	23	R	56	-69.7 %	-53	+143 %	+33	-26.3 %	-20		
		E	169					-86.4 %	-146			-66.9 %	-113		
CARIBBEAN INDEX SITES: <u>LACKING</u> Historical Data prior to 1985															
1901 Data Extrapolated using average 1901 to 1985 Population Trajectories from from Previous Table															
TOTAL: <u>LINEAR</u> & RAW DATA		L	815	R		247	R		1,385	-69.7 %	-568	+461 %	+1,138	+69.9 %	+570
TOTAL: <u>EXPONENTIAL</u> & RAW DATA		E	1,816			-86.4 %			-1,569	+461 %	+1,138	-23.7 %	-431		

ATL-Table 6 (c). South Western &amp; Eastern Atlantic sites

Index #	Index Sites <u>WITH</u> Historical Data	Population Estimate: mean # females/ yr			Estimated Change in Population						
					1901-1985		1985-2005		1901-2005		
		1901	1985	2005	%	# Fem	%	# Fem	%	# Fem	
ATLANTIC OCEAN: SOUTHWESTERN & EASTERN											
24	Brazil	R	2,338	L 827 E 434	R	468	-64.6 % -77.3 %	-1,507 -1,803	-43.3 % -11.7 %	-358 -62	-79.9 % -1,865
		L	360	L 99	L	0	-72.5 % -97.6 %	-261 -734	-100 % -77.8 %	-99 -14	-100 % -99.5 % -360 -748
25	Equatorial Guinea (Bioko)	L	360	L 99	L	0	-72.5 %	-261	-100 %	-99	-100 %
		E	752	E 18	E	4	-97.6 %	-734	-77.8 %	-14	-99.5 %
ATLANTIC OCEAN: SOUTHWESTERN & EASTERN											
TOTAL: <u>LINEAR</u> & RAW DATA		L	2,698	L 926	L	468	-65.7%	-1,768	-49.5%	-457	-82.7% -2,225
TOTAL: EXPONENTIAL & RAW DATA		E	3,090	E 452	E	472	-85.4%	-2,537	+4.4%	-76	-84.7% -2,623

**ATL-Table 6 (d). Regional Summaries**

Index #	Index Sites <u>WITH</u> Historical Data	Population Estimate: mean # females/ yr			Estimated Change in Population					
					1901-1985		1985-2005		1901-2005	
		1901	1985	2005	%	# Fem	%	# Fem	%	# Fem
ATLANTIC OCEAN: ENTIRE CARIBBEAN										
	TOTAL: <u>LINEAR</u> & RAW DATA	L 11,603	L 3,517	L 2,910	-69.7 %	-8,086	-17.3 %	-607	<u>-74.9 %</u>	-8,693
	TOTAL: <u>EXPONENTIAL</u> & RAW DATA	E 13,179	E 1,790	E 2,701	-86.4 %	-11,389	+50.9 %	+911	<u>-79.5 %</u>	-10,478
ENTIRE ATLANTIC OCEAN										
	TOTAL: <u>LINEAR</u> & RAW DATA	L 14,301	L 4,443	L 3,378	<u>-68.9 %</u>	-9,854	-24.0 %	-1,064	<u>-76.4 %</u>	-10,918
	TOTAL: <u>EXPONENTIAL</u> & RAW DATA	E 16,269	E 2,242	E 3,173	<u>-86.2 %</u>	-13,926	+41.5 %	+835	<u>-80.5 %</u>	-13,091

**ATL-Table 7.** Current population estimates and qualitative information about status and trends for reviewed hawksbill populations in the Atlantic Ocean. Population estimates are based on nesting females/yr, but where estimates are derived from numbers of nests, a bracketed figure of 3-5 nests per female is used to convert from numbers of nests to numbers of females, unless stated otherwise. Where the estimate is derived from total number of crawls, a conversion factor of 1.8 crawls per nest is used (based on Mortimer & Bresson, 1999).

Index #	Locality	Current Population Size	Comments (Source)	Status / Trends (Source)
<b>ATLANTIC OCEAN: INSULAR CARIBBEAN</b>				
14	Antigua / Barbuda	100-125 females/yr <sup>a</sup>          <sup>a</sup> 4 nests/female	Estimated 400-500 nests/yr (Meylan 1999). Most significant nesting site is Jumby Bay, Long Island, Antigua (Fuller <i>et al.</i> 1992, Richardson <i>et al.</i> 1999, Richardson <i>et al.</i> 2006). Shell exports to Japan since 1950: 3,146 turtles (4,216 kg).	Populations in Antigua and Barbuda are "remnants" (Fuller <i>et al.</i> 1992). No protection is afforded to ~35 additional Hawksbill beaches identified on Antigua and Barbuda by Groombridge and Luxmoore (1989), Joseph (1984), and Meylan (1983, 1999). No data are available to document current status at those sites. Pinchin Bay was the best site on Antigua 23 years ago (Meylan 1983). Number of nesting females at Jumby Bay, Long Island, stable during 1987-2001, and increased during 2002-2005 apparently in response to long-term protection (ATL-Table 2; Richardson <i>et al.</i> 2006).
15	Bahamas	100-333 females/yr	Estimated 500-1,000 nests/yr scattered throughout archipelago of over 700 islands and cays; no known nesting aggregations (K. Bjorndal, <i>in litt.</i> to J. Mortimer, 5 Nov 2006). The European shell trade was intense, and by the 1890s, average annual exports represented the shell of 3,122 turtles (4,186 kg) Assuming that half this shell came from nesters, McClenachan <i>et al.</i> (2006) estimated that an average of 1,561 nesters were taken annually during the 1890s; and, based on these data, identified Bahamas as one of the 7 major historic hawksbill nesting areas in the Caribbean. Shell exports to Japan since 1950: 14,876 turtles (19,934 kg)	Despite high Japanese demand in the 1960s and 1970s, export statistics indicate a decline of 82% in the average annual shell export from the 1890s to 1979. Carr <i>et al.</i> (1982) reported a considerable population decline in the 50 years prior to 1982. Hawksbills are protected by law, but nesting populations are threatened by poaching and unregulated coastal development (K. Bjorndal, <i>in litt.</i> to J. Mortimer, 5 Nov 2006; B. Riegl, pers. comm. to J. Mortimer, 9 Oct 2006).



16	<b>Barbados</b>	<b>~ 483 females/yr<sup>b</sup></b> <b><sup>b</sup> 4.1 nests/female (Beggs <i>et al.</i> 2007)</b>	Estimated ~1,981 nests/yr based on 2003-05 data (Beggs <i>et al.</i> 2007). In the 1960s and 1970s numbers and sizes of captured turtles decreased (Hunte 1984, Horrocks 1992). Shell exports to Japan since 1950: 2,401 turtles (3,218 kg).	Population seriously depleted by the mid-1980s, but increased significantly during 1997-2005, apparently in response to long-term protection and a moratorium in place since 1998 (Beggs <i>et al.</i> 2007).
	<b>Bonaire</b>	<b>8-14 females/yr</b>	Estimate based on 42 nests in 2006 (Nava and Uhr 2007, Dow and Eckert 2007).	Remnant population.
	<b>British Virgin Islands</b>	<b>No current estimates, but much reduced</b>	Recent study by McGowan <i>et al.</i> (MS in review) reports considerable numbers of foraging hawksbills thought to be derived mainly from major rookeries elsewhere in the Caribbean. Turtles can be legally killed in BVI during December-March. The BVI take may be impeding recovery of nesting populations on nearby St. John Island in the US-VI (Z. Hillis-Starr, <i>in litt.</i> to J. Mortimer 2006).	Historically the Hawksbill fishery was widespread. In the 1940s tortoiseshell was a major source of fishermen's income. Estimated catch has declined from 400 turtles in 1981, to 75 in 1985, to 32 in 1991 (Eckert <i>et al.</i> 1992). Recent studies (McGowan <i>et al.</i> MS in review) show nesting populations to have declined to critically low levels since the 1980s (Fletemeyer 1984), most probably due to historical exploitation.
17	<b>Cuba Doce Leguas Cays</b>	<b>400-833 females/yr</b>	Estimated 2,000-2,500 nests/yr. Full extent of nesting unknown; maximum total number of nests recorded on all 47 beaches in any one year for 1994-1998 was 409 (Moncada <i>et al.</i> , 1999). Current estimates of 2,000-2,500 nests/yr (Cuban Turtle Group <i>in litt.</i> to A. Abreu, Feb 2002), based on actual counts of 70 nests/yr. Legal take is currently 500 hawksbills per year (Carrillo <i>et al.</i> 1999). Shell exports to Japan since 1950: 106,948 turtles (170,047 kg) taken on foraging grounds.	Historical records indicate thousands of nesting females were captured annually during 19 <sup>th</sup> and 20 <sup>th</sup> centuries (Ballou 1888 as cited in McClenachan <i>et al.</i> 2006, McClenachan <i>et al.</i> 2006). In 1936 a closed season was introduced, and in 1961 government prohibited egg collection and disturbance of nesting females, suggesting concern about sustainability (Carrillo <i>et al.</i> 1999). Impact of current exploitation (500/yr) and current nesting trends are unknown, but suspected to be declining in some areas (Carrillo <i>et al.</i> 1999, Moncada <i>et al.</i> 1999), with small increases at other sites (Cuban Turtle Group to A. Abreu, Feb. 2002).

	<b>Dominican Republic</b>	<b>29-84 females / yr</b>	Estimate based on crawls at 11 sites: 3 beaches with <75-200 crawls (Ottenwalder, 1981), 7 sites totaling <150 crawls and 1 site with 25-100 crawls /yr in 2006 (Y. Leon pers. comm. to W. Dow, 2007; Dow & Eckert, 2007) .	Declining. Declines were underway in 1980s (Ottenwalder, 1981, 1987). Once considered a very important site for nesting hawksbills (Ottenwalder, 1981, 1987). Exploitation continues.
	<b>French West Indies Guadeloupean Archipelago</b>	<b>~ 40 - 66 females /yr</b>	Estimated ~200 nests / yr (Chevalier et al., 2005). Protective legislation for turtles implemented in 1991 greatly reduced number of turtles killed (Chevalier et al., 2005). Shell exports to Japan since 1950: 1572 turtles (2107 kg) from the French West Indies	Depleted & increasing. In mid-1980s, nesting levels very low (Meylan, 1999). Turtle populations increasing since 1991 (Chevalier et al., 2005; Chevalier et al., 2003), but deforestation threatens to feminize sex ratios of offspring produced (Kamel & Mrosovsky, 2006).
	<b>French West Indies Martinique</b>	<b>~50-100 females /yr</b>	In 1970s, Carr et al. (1982) considered exploitation of hawksbills in Martinique to be highest in Lesser Antilles; Lescure (1987) considered population "gravely threatened".	Dropsy (1987 cited in Meylan, 1999) estimated 245-375 nests (~50-125 females). Since 2004, conservation & research programmes implemented; 37 nesting hawksbills were tagged in 2006 (La Gazette des Karets, 2006).
	<b>Jamaica</b>	<b>200-275 females / yr</b>	Based on beach surveys from 1991-1996 (R. Kerr, pers. comm. to A. Meylan (2002).	More than 90% of coral reef habit destroyed since 1980 (R. Kerr, in litt. to A. Meylan 2002).
	<b>Grenada</b>	<b>&lt;25-56 females / yr</b>	Estimate based on <225-300 crawls per year (C. Lloyd and R. King, pers. comm.. to W. Dow 2007; M. Fastigipers. comm.. to W. Dow 2007; Dow and Eckert, 2007). Gazette et al. (2007) found evidence of decline in catch per unit effort for the in-water turtle fishery of Grenada.	Probably declining. Trend unknown but previous estimate of >500 females was not based on surveys (Meylan, 1999).
<b>18</b>	<b>Puerto Rico  Mona Island, Culebra Islands, Caja de Muertos, Humacao</b>	<b>Mona: 199-332 females / yr  Culebra: 8-13 females/yr  Caja de Muertos 13-22 females/y  Humacao: 30-50 females/y</b>	Estimated ~996 nests/ yr in 2001-2005 (Unpubl. data, R.P. van Dam & C.E. Diez; C.E. Diez in litt. to J. Mortimer 2006). Nesting activity at all four sites has increased during the survey periods: Caja de Muertos (1995-2003), +23%; Culebra Island (1993-2005), +190%; and Humacao (1987-2004), +930%. Shell exports to Japan since 1950: 4619 turtles (6190 kg) For the present assessment, Mona Island, with the longest history of monitoring, is the index site for Puerto Rico.	Populations appeared to be in decline until early 1990s (Unpubl. data, R.P. van Dam & C.E. Diez; C.E. Diez in litt. to J. Mortimer 2006).

	<b>St. Kitts and Nevis</b>	<p><b>St. Kitts:</b> <b>&lt;25-56 females/yr</b></p> <p><b>Nevis:</b> <b>&lt;43 females/yr</b></p>	<p>Estimate in St. Kitts based on &lt;225-300 crawls per year (K. Stewart pers. comm.. to W. Dow, 2007; Dow and Eckert, 2007) . Estimate in Nevis is based on &lt;25 crawls on each of 9 beaches and 25-100 crawls on the tenth (E. Pemberton pers. comm.. to W. Dow, 2007; Dow and Eckert, 2007)</p>	<p>Depleted population. Serious decline in recent decades (Eckert and Honebrink, 1992).</p>
	<b>Trinidad and Tobago</b>	<p><b>N. coast Trinidad:</b> <b>~150 females / yr</b></p>	<p>Estimate for N. coast Trinidad based on surveys conducted 2000-2004 (Livingstone, 2006). Significant but unquantified nesting reported for E. coast Trinidad and nearby Tobago (Livingstone, 2006).</p>	<p>Trends unknown. Current threats include: exploitation for shell (esp. Tobago); entanglement in gill nets and capture in shrimp trawls; and occasional take of nesting females for meat (Livingstone, 2006).</p>
<b>19</b>	<p><b>U.S. Virgin Islands</b></p> <p><b>Buck Island National Monument (St. Croix)</b></p>	<p><b>56 females / yr</b></p>	<p>Estimate based on long term monitoring data collected at Buck Island Reef National Monument (Z. Hillis-Starr, in litt. to J. Mortimer, 23 Oct 2006). Schmidt (1916) reported ongoing "wanton destruction" has led to population decline...and "every specimen, however small ...is landed and killed." Hawksbill eggs were also heavily exploited.</p>	<p>Hawksbill nesting is increasing at Buck Island National Park, with apparent spill over to beaches on adjacent St. Croix (Z. Hillis-Starr, in litt. to J. Mortimer, 23 Oct 2006).</p>
	<p><b>U.S. Virgin Islands</b></p> <p><b>Sites outside Buck Island National Monument</b></p>	<p><b>30-222 females / yr</b></p>	<p>Estimate based on 275-1,200 crawls per year (R. Boulon and S. Garner pers. comm.. to W. Dow, 2007; Dow and Eckert, 2007) St. Thomas has a long history of shell trade, and by 1914, hawksbill populations were considered much reduced from former years (Schmidt, 1916). In 1914, St. Thomas, St. John and their surrounding islets hosted the greatest number of hawksbills.</p>	<p>On St. Croix nesting is increasing at Sandy Point National Wildlife Refuge and also on several other beaches on SE coast being patrolled by conservation groups. But, similar increases have not been recorded at St. John, perhaps due to proximity to the legal turtle exploitation in British Virgin Islands (Z. Hillis-Starr, in litt., to J. Mortimer, 23 Oct 2006).</p>
<b>ATLANTIC OCEAN: WESTERN CARIBBEAN MAINLAND</b>				
	<p><b>Belize</b></p> <p>Manatee Bar, Sapodilla Cays, South Water Cay</p>	<p><b>~8-56 females / yr</b></p>	<p>Estimate based on 25-100 crawls per year in each of the three major areas of Manatee Bar, Sapodilla Cays, and South Water Cay (I. Majil pers. comm. to W. Dow, 2007; Dow and Eckert, 2007).</p>	<p>Declining. In the early 1990s, 100-150 nests (i.e., 20-50 females) were counted at Manatee Bar, and 30-40 nests (i.e., 6-13 females) in the southernmost cays (Smith et al., 1992). In the early 1900s, Belize supported a valuable tortoiseshell industry (Smith et al., 1992).</p>

	Colombia <b>Isla Fuerte</b>	<b>~19-93 females / yr</b>	Estimate based on 100-500 crawls per year (Ceballos-Fonseca, 2004)	Probably declining. Marked declines on offshore cays (Carr et al., 1982); many Caribbean sites have 25-100 crawls per year (Ceballos-Fonseca, 2004).
	Colombia <b>San Andres Archipelago</b>	<b>No current estimates, but much reduced</b>	Serrana, Serranilla, & Roncador, tiny sand cays 120-160 km NE of Providencia, likely source of much of 2,270 kg of shell exported by Cayman annually during 1932-1939 (Parsons, 1972). Shell exports to Japan since 1950: 767 turtles (1028 kg)	Despite its importance in the 1930s, these rookeries were almost extinct by 1981, nesting having declined significantly during the 1970s (Carr et al., 1982). In 1996, during a 7.5 month-long survey of archipelago nesting beaches, only 21 hawksbill nests were seen (Cordoba et al., 1998).
20	Costa Rica <b>Tortuguero National Park</b>	<b>&lt; 10 females /yr</b>	Hawksbills have been protected at Tortuguero for decades. The continued population decline may be due to legal & illegal, directed & incidental take where the reproductive animals forage (Troëng et al., 2005), & possibly low clutch survival (Harrison et al., 2003). Shell export to Japan since 1950: 6717 turtles (9001 kg)	Trend analyses indicate nesting declined 77.2-94.5% between 1956 and 2003 (Troëng et al., 2005).
	Costa Rica <b>Cahuita National Park</b>	5-9 females/yr	Estimate from Hancock, 2007.	Trend unknown, but suspected to be in decline for the same reasons hawksbills are declining at Tortuguero.
	Honduras <b>Bay Islands</b>	<10 females / yr	Aerial & ground surveys during 1982-1987 revealed only sparse nesting (Cruz & Espinal, 1987). Average of 22 nests / yr recorded by monitoring Archipelago of Cayos Cochinos in 1999 and 2000 (Aronne, 2000a, 2000b). Shell exports to Japan since 1950: 7507 turtles (10,059 kg)	Remnant population. A major hawksbill rookery in the 16 <sup>th</sup> & 17 <sup>th</sup> centuries (McClenachan et al., 2006). 20 <sup>th</sup> Century declines have been significant (Carr et al., 1982; Meylan, 1999).

21	<i>Mexico Yucatán Peninsula</i>	2,672 nests/ yr  534 - 891 females /yr	Estimate based on average for 2001-2006 (A. Abreu-Grobois in litt. to J. Mortimer, 9 Feb 2007). From 1977-2005 population variation mirrored across all three states of the Yucatán Peninsula (A. Abreu-Grobois et al., 2005). Shell exports to Japan since 1950: 1696 turtles (2273 kg)	The Yucatán Peninsula once hosted the best fishing for caret in the Americas (Parsons, 1972). Yucatán nesting hawksbills believed to have declined prior to 1977. Nesting increased dramatically between 1977 and 1999, followed by significant declines between 1999 and 2004, and stabilization between 2004 & 2005 (Abreu-Grobois, pers. comm. to M. Donnelly, 2006). Reported nestings in 2007 only 50% of 1999 levels (E. Cuevas, in litt. 28 Aug 2007). Mexican researchers suspect recent declines due to extraction at low levels and/or impacts on marine habitats (Abreu-Grobois et al., 2005).
22	<i>Nicaragua El Cocal</i>	15-25 females / yr	Estimated ~75 nests/yr during 2000 (Lagueux & Campbell, 2005). Almost all eggs collected annually, nesting females killed on the beach & entangled in commercial fishing gear (C. Lagueux, in litt. to J. Mortimer, 2001).	Based on beach surveys and interviews, researchers conclude declines of >75% since the 1970s (Lagueux & Campbell, 2005).
	<i>Nicaragua Miskito Coast</i>	unknown numbers of foraging turtles	Hawksbill exploitation was year-round, estimated at ~ 1,000-1,200 turtles /yr (Nietschmann, 1981). Lobster divers captured hawksbills whenever encountered, and in 1992 reported them as becoming rare (J. Mortimer, unpub. data from Miskito Coast Protected Area Project of the Caribbean Conservation Corporation). Shell exports to Japan from Nicaragua since 1950: 11,779 turtles (15,784 kg)	Decline in foraging hawksbills of >92% in 28 yrs (Lagueux, 1998). In Tasbabaune community, number of hawksbills killed semi-annually dropped from 67 in 1968 & 1971 (Nietschman, 1972, 1973), to only 14 during 1995-1997 (Lagueux, 1998).
	<i>Nicaragua Pearl Cays</i>	30 - 52 females / yr	Estimated ~155 nests/yr during 2000-2002 (Lagueux et al., 2003), 176 in 2004, 205 in 2005, and 211 in 2006 (C. Campbell, pers. comm to J. Mortimer, 23 Feb 2007). In 1971-72, 90-95% of nests excavated by fishermen (Nietschmann, cited in Groombridge & Luxmoore, 1989). Lagueux et al. (2003) report ~100% egg collection & many nesting females killed prior to 2000.	Numerous interviews indicate population decline (C. Lagueux, in litt. to J. Mortimer, 2001). Since 2000, exploitation reduced by community awareness campaign, but coastal development by foreign nationals poses extreme threat to nesting habitat (Lagueux et al., 2003).

23	<i>Panamá Bastimentos Island National Marine Park</i>	27 - 45 females / yr	<p>Estimated ~136 nests / yr during 2003-2005 (Meylan et al., 2006).</p> <p>Protection was implemented in 1988. Lack of mammalian predators and artificial lighting, and limited human presence have provided favourable nesting conditions (Meylan et al., 2006).</p> <p>Shell exports to Japan since 1950: 150,863 turtles (202,157 kg)</p>	Nesting activity increased between 1990 and 2005, apparently in response to protection. Nests recorded at Small Zapatilla Cay during first 3 weeks of July were: for 1991, 1993, 1997, mean = 4.3; for 2000, 2002-2005, mean = 13.0 (Meylan et al., 2006).
	<i>Panamá Chiriqui Beach</i>	84 - 140 females / yr	<p>Estimated ~421 nests / yr during 2003-2005 (Meylan et al., 2006)</p> <p>Playa Chiriqui, historically the most significant rookery in the region, is severely depleted. Recently gained protected status as Damani-Guariviara Wetland; but, threats from poaching &amp; predators (esp. dogs), are difficult to address on this mainland beach (Meylan et al., 2006).</p> <p>Diez et al. (2002) recorded unusually low hawksbill numbers in optimal foraging habitat in the Kuna Yala Archipelago (Panamá), and attributed this to over-exploitation of nearby rookeries.</p>	Nesting population has declined by > 95% during the past 50 years (Carr, 1956; Carr et al., 1982; Meylan & Donnelly, 1999; Ordoñez, pers. comm. to A. Meylan).
	<i>Venezuela Los Roques &amp; Paria region</i>	~ 32 - 53 females / yr	<p>Estimated ~159 nests / yr (H. Guada, in litt. to J. Mortimer, 2006).</p> <p>Serious threats include illegal take, destruction of foraging &amp; nesting habitats, &amp; incidental capture in fishing gear (Buitrago &amp; Guada, in final review cited in H. Guada in litt. to J. Mortimer, 2006).</p> <p>Significant domestic trade in shell for handicrafts and spurs for cock fighting continues (H. Guada, in litt. to J. Mortimer, 2006).</p> <p>Shell exports to Japan since 1950: 2349 turtles (3148 kg)</p>	Hawksbill nesting occurs on the continental coastline (> 55 nests/ yr) as well as on the islands, especially Los Roques (~ 104 nests/yr) (Buitrago & Guada, in final review, cited in H. Guada in litt. to J. Mortimer, 2006). But populations are much reduced primarily due to massive exploitation for shell in the 1960s and 1970s.




# ATLANTIC OCEAN: SOUTH WESTERN

24	<b>Brazil</b>	<b>350-585 females / yr</b>	<p>Estimated 1,750 nests / yr (N. Marcovaldi, <i>in litt.</i> to J. Mortimer, 17 Oct 2006).</p> <p>Nesting once extended from north Rio de Janeiro State all the way to the Ceará State (N. Marcovaldi, <i>in litt.</i> to J. Mortimer, 2001), but is today restricted primarily to northern Bahia and Sergipe (~1300 nests annually), Rio Grande do Norte, near Pipa (~450 nests in 2002-03), and only scattered nesting elsewhere (Marcovaldi, 2005).</p> <p>Numerous cases of viable hybrid hatchlings (crosses between hawksbills &amp; loggerheads) and nesting hybrid females recorded during past ten years (Lara-Ruiz et al. 2006). Bass <i>et al</i> (1996) and Bowen et al., 2007 reported other hybrids.</p> <p>Shell exports to Japan since 1950: 11 turtles (15 kg)</p>	<p>More than 80% population decline during the past 105 years extrapolated from reduced nesting distribution (N. Marcovaldi, <i>in litt.</i> to J. Mortimer, 2001), directed take of females &amp; eggs, manufacture of shell ornaments, incidental capture in fishing gear, &amp; habitat destruction before 1982. Since protection in 1982, the decline in the nesting population has stopped; studies from 1991 to 2006 on the population in northern Bahia and Sergipe show an increasing trend in nest numbers (Marcovaldi et al., <i>in press</i>). Hybridization may be a threat.</p>
----	---------------	-----------------------------	--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

# ATLANTIC OCEAN: EASTERN

25	<b>Equatorial Guinea Bioko</b>	<b>&lt; 7 females / yr</b>	<p>Comprehensive surveys recorded 13 females in 1996-97 and 1 in 1997-98 (Tomás et al., 2000). During 2001-05 only 4-7 nests recorded each year (Rader et al., 2006).</p> <p>Greens and hawksbills nest on 20 km of beach on southern Bioko. In 1940s, 200-300 nesters (of all species) taken daily at peak season; down to 50-100 in mid-1980s (T. Butynski <i>in litt.</i> to K. Bjørndal, 20 April 1986 cited in Groombridge &amp; Luxmoore, 1989). Hawksbills intensely exploited for eggs and shell (Castroviejo et al., 1994; Graff, 1996).</p>	<p>Population declining (Fretey &amp; Formia, <i>in litt.</i> to J. Mortimer, 2001; A. Formia, <i>in litt.</i> 28 Aug 2007).</p>
	<b>São Tomé and Príncipe</b>	<b>14 - 27 nesting females / yr</b>	<p>Estimated ~50 nests in São Tomé and ~20-30 nests in Príncipe (1998-2001). Approximately 80% of nesting females and eggs collected annually (Dontaine, <i>in litt.</i> to J. Mortimer, 2001.) Over-exploitation for tortoiseshell trade (J. Fretey, <i>in litt.</i> to J. Mortimer, 2002).</p>	<p>Population declining (Fretey &amp; Formia, <i>in litt.</i> to J. Mortimer, 2001; Dontaine, <i>in litt.</i> to J. Mortimer, 2001)</p>



**ATL- Figure 1.** Recorded hawksbill nesting in the Yucatan Peninsula 1977-2005 (Source: Abreu-Grobois *et al.*, 2007[Myrtle Beach presentation] o Resultados Mesa de Tendencias- XIV Taller Regional de Programas de Investigación y Manejo de Tortugas Marinas en la Península de Yucatán, Parque X'Caret, Quintana Roo, México, 8-10 Noviembre, 2006).